A SHARP UPPER BOUND FOR SAMPLING NUMBERS IN L_2

MATTHIEU DOLBEAULT[1], DAVID KRIEG[2,3], AND MARIO ULLRICH[2]

Abstract. For a class F of complex-valued functions on a set D, we denote by $g_n(F)$ its sampling numbers, i.e., the minimal worst-case error on F, measured in L_2, that can be achieved with a recovery algorithm based on n function evaluations. We prove that there is a universal constant $c \in \mathbb{N}$ such that, if F is the unit ball of a separable reproducing kernel Hilbert space, then

$$g_{cn}(F)^2 \leq \frac{1}{n} \sum_{k \geq n} d_k(F)^2,$$

where $d_k(F)$ are the Kolmogorov widths (or approximation numbers) of F in L_2. We also obtain similar upper bounds for more general classes F, including all compact subsets of the space of continuous functions on a bounded domain $D \subset \mathbb{R}^d$, and show that these bounds are sharp by providing examples where the converse inequality holds up to a constant. The results rely on the solution to the Kadison-Singer problem, which we extend to the subsampling of a sum of infinite rank-one matrices.

1. Introduction and main results

The general question of how well point-wise evaluations perform for approximating a function, which is often called *sampling recovery* or approximation using *standard information*, is a classical question in theoretical and applied mathematics. A historical treatment and various basics may be found in the monographs [9, 10, 11, 49, 61] for general approximation theory and in [41, 42, 43] for information-based complexity. It is of particular interest to compare the *power of function evaluations* with the power of optimal linear measurements (which

[1]Sorbonne Université, Paris, France
[2]Institut für Analysis, Johannes Kepler Universität Linz, Austria
[3]RICAM, Austrian Academy of Sciences, Linz, Austria.
E-mail address: matthieu.dolbeault@sorbonne-universite.fr, david.krieg@jku.at, mario.ullrich@jku.at

2010 *Mathematics Subject Classification.* 41A25, 41A45, 46B09, 46B15, 60B20.
Key words and phrases. L_2-approximation, information-based complexity, least squares, rate of convergence, random matrices, Kadison-Singer.
DK is supported by the Austrian Science Fund Project F5506, which is part of the Special Research Program *Quasi-Monte Carlo Methods: Theory and Applications.*

could be Fourier coefficients or derivatives), since the latter are well understood in many cases and easier to handle from a theoretical point of view, while the first are of larger practical relevance. The quest for a systematic comparison has attracted much attention recently. We will describe the history and related results below after presenting the setting and the main results, see also Section 1.1.

The *power* of a given class of measurements is often expressed in terms of the minimal error achievable with a given amount of such information. Here, we consider L_2-approximation in a worst-case setting, so that these minimal errors correspond to sampling numbers and Kolmogorov (or approximation) numbers, as we summarize below.

Let $(D, \mathcal{A}, \mu)$ be a measure space and $L_2 := L_2(D, \mathcal{A}, \mu)$ be the space of square-integrable complex-valued functions on D. Let F be a set of functions contained in L_2. The *Kolmogorov widths* of F in L_2 are defined by

$$d_k(F) := \inf_{\substack{\ell_1,\dots,\ell_k\colon F\to\mathbb{C} \\ \varphi_1,\dots,\varphi_k\in L_2}} \sup_{f\in F} \left\| f - \sum_{i=1}^{k} \ell_i(f)\,\varphi_i \right\|_{L_2}.$$

This is the worst-case error of an optimal approximation within a linear space of dimension k. It coincides with the kth approximation number (or linear width) of F, which is the worst-case error of an optimal linear algorithm that uses at most k linear functionals as information, see Remark 5. On the other hand, the *sampling numbers* are given by

$$g_n(F) := \inf_{\substack{x_1,\dots,x_n\in D \\ \varphi_1,\dots,\varphi_n\in L_2}} \sup_{f\in F} \left\| f - \sum_{i=1}^{n} f(x_i)\,\varphi_i \right\|_{L_2},$$

i.e., $g_n(F)$ is the minimal worst-case error of linear algorithms based on n function evaluations. Therefore, the task is to compare the numbers $d_k(F)$ and $g_n(F)$.

It is clear that we have $g_n(F) \geq d_n(F)$. Here, we aim for an upper bound of $g_n(F)$ in terms of the numbers $d_k(F)$. We first describe the situation where F is the unit ball of a separable reproducing kernel Hilbert space (RKHS). A priori, it is not clear whether such a bound is even possible. And indeed, there can be no such bound in the case that $(d_k(F)) \notin \ell_2$. More precisely, it is shown in [13] that for any non-negative and non-increasing sequence $(\sigma_k) \notin \ell_2$ and any sequence (τ_n) tending to infinity, e.g. $\tau_n = \log\log n$, there exists a RKHS with unit ball F such that $d_k(F) = \sigma_k$ for all k but $\limsup_{n\to\infty} \tau_n \cdot g_n(F) > 0$.

The situation is completely different when $(d_k(F)) \in \ell_2$, which is equivalent to assuming that the kernel K of the Hilbert space has finite trace

$$\int_D K(x,x)\,d\mu(x) < \infty, \tag{1}$$

see, e.g., [37]. Under this assumption, first upper bounds on $g_n(F)$ in terms of the numbers $d_k(F)$ were obtained more than 20 years ago in [59]. These upper bounds were later improved in [27, 30, 38]. On the other hand, a lower bound from [17, Theorem 2] tells us how far these improvements might go: for every non-negative and non-increasing $(\sigma_k) \in \ell_2$, there exists a separable RKHS with unit ball F such that $d_k(F) = \sigma_k$ for all $k \in \mathbb{N}$ and

$$g_{\lfloor m/8 \rfloor}(F) \;\geq\; \sqrt{\frac{1}{m} \sum_{k \geq m} d_k(F)^2} \tag{2}$$

for infinitely many values of $m \in \mathbb{N}$. And indeed, it turns out that this is already the worst possible scenario. The main result of this paper is an upper bound, which matches the above lower bound (2) up to a universal constant, and which is true for any separable reproducing kernel Hilbert space.

Theorem 1. *There is a universal constant $c \in \mathbb{N}$ such that the following holds. Let μ be a measure on a set D and let $F \subset L_2(\mu)$ be the unit ball of a separable RKHS on D such that the finite trace assumption* (1) *holds. Then, for all $m \in \mathbb{N}$, we have*

$$g_{cm}(F) \;\leq\; \sqrt{\frac{1}{m} \sum_{k \geq m} d_k(F)^2}.$$

This settles the question on the power of standard information compared to general linear information for the problem of L_2-approximation on Hilbert spaces, and solves the open problems from [17, 27], Open Problem 140 in [43], as well as Outstanding Open Problem 1.4 in [11] for L_2-approximation. The latter is discussed in Example 28, where we consider tensor product spaces. We note that the case of L_p-approximation ($p \neq 2$) is widely open. A slightly stronger version of Theorem 1 and explicit constants are given in Theorem 23.

Let us add that, in principle, Theorem 1 does only imply the *existence* of (linear) sampling algorithms achieving the error bound. However, all upper bounds on $g_n(F)$ will be obtained by a suitable (unregularized) *least squares method*, see Remark 7 and Section 5.

Theorem 1 is a direct continuation of the series of works initiated in [27], in which the sampling numbers were bounded by

$$g_{\lfloor c\, m \log m \rfloor}(F) \;\leq\; \sqrt{\frac{1}{m} \sum_{k \geq m} d_k(F)^2},$$

see also [19, 55], and an improvement from [38], where the logarithmic oversampling was removed in exchange for an additional factor $\sqrt{\log m}$ on the right hand side.

The ingredients for the proof are still the existence of good point sets with $\mathcal{O}(m \log m)$ points from [27], and a subsampling of $\mathcal{O}(m)$ points based on the solution to the Kadison-Singer problem [34]. The Kadison-Singer subsampling has already been applied for the related problem of sampling discretization in [32] (see [20] for a survey) and was subsequently introduced to the study of sampling numbers in [38, 50]. In these papers, the subsampling was, roughly speaking, only performed for a finite-dimensional sub-problem which resulted in the excessive factor $\sqrt{\log m}$ in [38]. The new ingredient here is an infinite-dimensional version of the subsampling theorem that might be of independent interest, see Proposition 17.

If we apply Theorem 1 and the lower bound from [17] to sequences with polynomial decay, we obtain the following characterization.

Corollary 2. *Let F be the unit ball of a separable RKHS with*

$$d_n(F) \lesssim n^{-\alpha} \log^{-\beta} n \tag{3}$$

for some $\alpha \geq 1/2$, $\beta \in \mathbb{R}$ and $c > 0$. Then

$$g_n(F) \lesssim \begin{cases} n^{-\alpha} \log^{-\beta} n & \text{if } \alpha > 1/2, \\ n^{-\alpha} \log^{-\beta+1/2} n & \text{if } \alpha = 1/2 \text{ and } \beta > 1/2. \end{cases} \tag{4}$$

Moreover, there exist classes F such that these bounds are sharp.

Here, $a_n \lesssim b_n$ means that there is a constant $c > 0$ such that $a_n \leq c b_n$ for all but finitely many $n \in \mathbb{N}$; later we will also use the symbols $\gtrsim$ and $\asymp$ which are defined accordingly. It is clear from Theorem 1 that the hidden constant in (4) is given by the product of the hidden constant in (3) and a constant that only depends on α and β.

We now turn to general function classes F that are assumed to satisfy the following assumption.

Assumption A. Let F be a class of complex-valued functions on a set D and let μ be a measure on D. We say that F and μ satisfy Assumption A, if there is a metric on F such that F is continuously embedded into L_2, separable, and function evaluation $f \mapsto f(x)$ is, for each $x \in D$, continuous on F.

Note that Assumption A is satisfied, for example, if

- F is a separable subset of the space of bounded functions equipped with the maximum distance and the measure μ is finite, **or**
- F is the unit ball of a separable normed space that is continuously embedded in L_2 and on which function evaluation at each point is a continuous functional, **or**
- F is a countable set of square-integrable functions, equipped with the discrete metric.

In this setting, we prove the following bound.

Theorem 3. *Let $0 < p < 2$. There is a constant $c_p \in \mathbb{N}$, depending only on p, such that for any F and μ that satisfy Assumption A and all $m \in \mathbb{N}$,*

$$g_{c_p m}(F) \,\leq\, \left(\frac{1}{m}\sum_{k\geq m} d_k(F)^p\right)^{1/p}.$$

Theorem 3 is an improvement over [28], where again we removed the excessive logarithmic factor. We will also show that the result is not true for $p = 2$, see Example 30. However, we provide a variant of Theorem 3 under the weaker condition $\big((\log k)^s d_k(F)\big) \in \ell_2$ for some $s > 1/2$ in Section 6.2. This leads to the following corollary.

Corollary 4. *Let F and μ satisfy Assumption A and*

$$d_n(F) \,\lesssim\, n^{-\alpha}\log^{-\beta} n \tag{5}$$

for some $\alpha > 0$ and $\beta \in \mathbb{R}$. Then

$$g_n(F) \,\lesssim\, \begin{cases} n^{-\alpha}\log^{-\beta} n & \text{if } \alpha > 1/2, \\ n^{-\alpha}\log^{-\beta+1} n & \text{if } \alpha = 1/2 \text{ and } \beta > 1, \\ 1 & \text{otherwise.} \end{cases} \tag{6}$$

Moreover, there exist classes F such that these bounds are sharp.

Again, the hidden constant in (6) is given by the product of the hidden constant in (5) and a constant that only depends on α and β. The difference compared to unit balls of RKHSs is the case $\alpha = 1/2$, where we need $\beta > 1$ instead of $\beta > 1/2$ and lose a factor $\log n$ instead of $\sqrt{\log n}$, see Example 29. In addition, if $(d_k(F)) \notin \ell_2$, then $g_n(F)$ might be bounded below by a constant, opposite to the RKHS setting where $g_n(F)$ tends to zero as soon as $d_k(F)$ does, see [13]. However, for $\alpha > 1/2$, the results for general classes are just as strong as before.

1.1. **Remarks and related literature.**

We want to add several remarks on the history of the result and related topics.

Remark 5 (Equivalent widths)**.** There are several quantities to measure the "width" of a set F. Although we work here with the Kolmogorov numbers $d_k(F)$ as benchmark, let us add that these quantities coincide in L_2 with the *approximation numbers* of F, i.e.

$$d_k(F) \,=\, a_k(F) \,:=\, \inf_{\substack{\ell_1,\dots,\ell_k\colon F\to\mathbb{C}\text{ linear}\\ \varphi_1,\dots,\varphi_k\in L_2}} \sup_{f\in F} \Big\| f - \sum_{i=1}^{k} \ell_i(f)\,\varphi_i \Big\|_{L_2},$$

as the infimum in the definition of $d_k(F)$ for given $\varphi_1, \dots, \varphi_k$ is attained by the L_2-orthogonal projection onto their span, which is linear in any case. The approximation numbers of a class represent the worst-case

error of an optimal linear algorithm that uses at most k linear functionals as information. If F is the unit ball of some Hilbert space H, then the approximation numbers agree with the *singular values* of the identity $\mathrm{Id}\colon H \to L_2$. In this case, the $d_k(F)$ also coincide with the *Gelfand k-widths* $c_k(F)$, which represent the minimal worst-case error of (possibly non-linear) algorithms based on k arbitrary linear functionals, see, e.g., Chapter 4 in [41].

Remark 6 (Extreme classes F)**.** It is interesting to note that the lower bound (2) from [17] is attained already for univariate Sobolev spaces of periodic functions. By Theorem 1, this means that these basic classes already represent the most difficult RKHSs for sampling recovery when the numbers $d_k(F)$ are fixed.

Remark 7 (Least squares methods)**.** The upper bounds in Theorem 1 and 3 are proven for a weighted least squares algorithm using samples from a set of cm points that is subsampled from a set of $cm \log m$ i.i.d. random points, see Section 5. Depending on the function class F, the algorithm using the full set of random points may be constructive but the subsampling is based on an existence result from [34] and is therefore not constructive. It would be very interesting to make the subsampling constructive, see Remark 21.

Remark 8 (Spline algorithm)**.** Let F be the unit ball of a RKHS H. If we fix the sampling points $x_1, ..., x_n$, it is known that the smallest possible worst case error is achieved by the spline algorithm

$$S_n(f) := \operatorname*{argmin}_{g \in H\colon\, g(x_i)=f(x_i)} \|g\|_H,$$

that is,

$$\inf_{\varphi_1,\dots,\varphi_n \in L_2} \sup_{f\in F} \Big\| f - \sum_{i=1}^n f(x_i)\, \varphi_i \Big\|_{L_2} = \sup_{f\in F} \Big\| f - S_n(f) \Big\|_{L_2},$$

see e.g. [54, Theorem 5.1]. The function $S_n(f)$ is also known as the minimal norm interpolant and, by the famous *representer theorem*, can be expressed as a linear combination of the kernel functions $K(x_i, \cdot)$, see e.g. [57, Proposition 12.32]. Therefore, our upper bounds are true not only for the least squares algorithm, but also for the kernel-based approximation $S_n(f)$. Both types of algorithms are common in the context of learning, see e.g. the seminal paper [9].

Remark 9 (The power of i.i.d. sampling)**.** It is remarkable that, up to a logarithmic factor, the upper bound from Theorem 1 is achieved with high probability for i.i.d. random sampling points, see [27, 55]. In regard of the personal history of the authors DK and MU, Theorem 1

is a byproduct of a series of work on the power of i.i.d. sampling for approximation and integration problems that started in [14, 15] and was also continued in [18, 23, 25, 26].

Remark 10 (Expected error)**.** A different approach to L_2-approximation is by using randomized algorithms and taking the worst case expected error instead of a worst case deterministic error. The results in this randomized setting are quite different; the error of optimal algorithms does not depend on the tail of the sequence $(d_k(F))$. We refer to [7, 8, 21, 33, 43, 60].

Remark 11 (Upper bounds for infinite trace)**.** We note that our bounds make sense also if $d_k(F)$ is infinite for small k, but they are useless if the *tail* of $(d_k(F))$ is not square-summable, which is the case, e.g., if F is the unit ball of a RKHS with infinite trace, see (1).
An alternative approach is to bound the numbers $g_n(F)$ by the Kolmogorov widths $d_k(F, L_\infty)$ in L_∞: it is shown in [50] that there is a universal constant $c \in \mathbb{N}$ such that $g_{cm}(F) \leq c\, d_m(F, L_\infty)$ for probability spaces $(D, \mathcal{A}, \mu)$. Although this bound is sometimes weaker than Theorem 3 (see Example 1 in [28]), it has the great advantage that it may be applied in situations where the Kolmogorov widths in L_2 are not square-summable, see, e.g., [52, 53]. It would be very interesting to see whether it is possible to unify the two approaches.

Remark 12 (Tractability)**.** Assume now that a whole sequence of classes F_d is given, where d could be the dimension of the underlying domain. For some classes we know that the curse of dimensionality is present, if only standard information (function values) is allowed, while the problem is tractable for general linear information, see, e.g., [16, 44, 56]. However, since the constants from Theorems 1 and 3 are independent of the dimension, it is possible to transfer certain tractability properties from linear information to standard information, see, e.g., [19, 24, 43].

Remark 13 (Separability of F)**.** Contrarily to the ℓ_2-summability of the Kolmogorov widths, it should be possible to remove the separability assumption on the class F, at least in Theorem 1, by adding a term $\mathrm{tr}_0(K)/m$ inside the square root in the right-hand side, as done in [37].

Remark 14 (Discretization of continuous frames)**.** A related problem is the question whether a *continuous frame* for a Hilbert space may be sampled to obtain a frame, see [6] for details. This problem, which was originally posed in the physics book [1], has only recently been solved in [12], see also the survey [5]. Although seemingly independent, this line of research uses remarkably similar methods. We leave it to future research to better understand and expand the connections.

1.2. **Outline.** The rest of the paper can be outlined as follows. Sections 2–5 form the proof of Theorem 1. In Section 2, we collect some basics on the RKHS setting. In Section 3, we obtain our initial sample of $\mathcal{O}(m \log m)$ points based on a concentration inequality for infinite matrices. The subsampling is performed in Section 4, which applies the solution to the Kadison-Singer problem in a slightly original way, leading to the core of the proof in Section 5. In Section 6, we prove our results for general function classes by constructing a suitable RKHS, on which a local version of Theorem 1 (Theorem 23) can be applied. Finally, in Section 7, we present examples, applying our result to tensor product problems and showing that our upper bounds are sharp.

Acknowledgement: The authors thank Albert Cohen, Daniel Freeman, Aicke Hinrichs and Erich Novak for useful discussions and detailed corrections. Moreover, DK and MU want to acknowledge that MD obtained Theorem 1 on his own and presented it at the *MASCOT-NUM Workshop on "Optimal Sampling for Approximation"* at the IHP in Paris on March 10, 2022. The current work is the product of subsequent collaboration.

2. Hilbert space setting

We first consider the case where F is the unit ball of a separable Hilbert space H with reproducing kernel $K \in \mathbb{C}^{D \times D}$. We refer to [37] and references therein for theoretical background on RKHSs.

Thanks to the finite trace assumption (1), we know that the identity map $\mathrm{Id}\colon H \to L_2$ is Hilbert-Schmidt, thus its left and right singular vectors $(b_k)_{k\in\mathbb{I}}$ and $(\sigma_k b_k)_{k\in\mathbb{I}}$ are orthonormal families in L_2 and H, respectively. Here, we only list the singular vectors with respect to the nonzero singular values $\sigma_k > 0$, and the index set is of the form $\mathbb{I} = \{k \in \mathbb{N}_0 \colon k < M\}$ with $M \in \mathbb{N} \cup \{\infty\}$. The singular vectors satisfy

$$\langle f, b_k \rangle_{L_2} = \langle f, \sigma_k^2 b_k \rangle_H \quad \text{for all } f \in H \text{ and } k \in \mathbb{I}.$$

We use the convention that $\mathbb{N}_0 := \{0, 1, 2, \dots\}$ and the singular values are arranged in a non-increasing order. In particular, $\sum_{k\in\mathbb{I}} \sigma_k^2 < \infty$ and the Kolmogorov width $d_m(F) = \sigma_m$ is attained by the L_2-orthogonal projection P_m onto $V_m = \operatorname{span}\{b_k \colon k < m\}$. Moreover, the separability of H ensures that the equality

$$K(x,y) = \sum_{k\in\mathbb{I}} \sigma_k^2 \, b_k(x)\overline{b_k(y)}$$

holds for all $x, y \in D_0$ with some set $D_0 \subset D$ satisfying $\mu(D \setminus D_0) = 0$. We therefore have the identity

$$f(x) = \sum_{k\in\mathbb{I}} \langle f, b_k \rangle_{L_2} \, b_k(x) \quad \text{for all } f \in H \text{ and } x \in D_0. \tag{7}$$

Our sampling points will be contained in the set D_0.

As a consequence of the following lemma, we only have to show the validity of Theorem 1 for all $1 \le m < M$.

Lemma 15. *Let $M = \min\{m \in \mathbb{N}\colon d_m(F) = 0\} < \infty$. Then we have $g_n(F) = 0$ for all $n \ge M$.*

Proof. For $x \in D_0$, we write $b(x) = (b_0(x), \dots, b_{M-1}(x))$. Then there are points $x_0, \dots, x_{M-1} \in D_0$ such that every $b(x)$ is contained in the span of the vectors $b(x_i)$. We write $b(x) = \sum \varphi_i(x) b(x_i)$ with coefficients $\varphi_i(x) \in \mathbb{C}$. By (7), we have

$$f(x) = \sum_{k<M} \langle f, b_k\rangle_{L_2} \sum_{i<M} \varphi_i(x) b_k(x_i) = \sum_{i<M} f(x_i)\varphi_i(x),$$

for all $x \in D_0$ and $f \in H$. Thus, the identity $f = \sum f(x_i)\varphi_i$ holds almost everywhere. Moreover, the functions $b_0, \dots, b_{M-1}$ restricted to D_0 form a basis of $\operatorname{span}\{\varphi_i\colon i < M\}$, and thus $\varphi_i \in L_2$. □

We fix an integer $1 \le m < M$ for the rest of the proof of Theorem 1.

3. Concentration inequality

As proposed in [27] and applied in [19, 28, 37, 38, 55], we define the probability density

$$\rho_m(x) = \frac{1}{2}\left(\frac{1}{m}\sum_{k<m} |b_k(x)|^2 + \frac{\sum_{k\ge m} \sigma_k^2 \, |b_k(x)|^2}{\sum_{k\ge m} \sigma_k^2}\right).$$

and draw i.i.d. random points $x_1, \dots, x_n \in D$ according to this density. We define the M-dimensional vectors $y_1, \dots, y_n$ by

$$(y_i)_k = \begin{cases} \rho_m(x_i)^{-1/2}\, b_k(x_i) & \text{if } 0 \le k < m, \\ \rho_m(x_i)^{-1/2}\, \gamma_m^{-1}\, \sigma_k\, b_k(x_i) & \text{if } m \le k < M, \end{cases}$$

where

$$\gamma_m := \max\left\{\sigma_m, \sqrt{\frac{1}{m}\sum_{k\ge m} \sigma_k^2}\right\} > 0.$$

Note that $\rho_m(x_i) > 0$ almost surely. It follows from these definitions that $y_i \in \ell_2(\mathbb{I})$ with

$$\|y_i\|_2^2 = \rho_m(x_i)^{-1}\left(\sum_{k<m} |b_k(x_i)|^2 + \gamma_m^{-2}\sum_{k\ge m} \sigma_k^2\, |b_k(x_i)|^2\right) \le 2m,$$

and

$$\mathbb{E}(y_i y_i^*) = \operatorname{diag}(1, \dots, 1, \sigma_m^2/\gamma_m^2, \sigma_{m+1}^2/\gamma_m^2, \dots) =: E,$$

with $\|E\|_{2\to 2} = 1$ since $\sigma_k^2/\gamma_m^2 \le 1$ for $k \ge m$. Here, $\operatorname{diag}(v)$ denotes a diagonal matrix with diagonal v, and $\|\cdot\|_{2\to 2}$ denotes the spectral norm of a matrix.

We apply the following concentration inequality for infinite matrices, which was proved by Mendelson and Pajor in [35, Theorem 2.1]. We use a version of this result from [37, Theorem 1.1] and [38, Theorem 5.3].

Lemma 16. *Let $n \geq 3$ and $y_1, \dots, y_n$ be i.i.d. random sequences from $\ell_2(\mathbb{I})$ satisfying $\|y_i\|_2^2 \leq 2m$ almost surely and $\|E\|_{2\to 2} \leq 1$, with $E = \mathbb{E}(y_i y_i^*)$. Then, for $0 \leq t \leq 1$,*

$$\mathbb{P}\left(\left\| \frac{1}{n} \sum_{i=1}^{n} y_i y_i^* - E \right\|_{2\to 2} > t \right) \leq 2^{3/4}\, n \, \exp\left(-\frac{nt^2}{42\, m} \right).$$

For $t = 1/2$, this probability is less than $1/2$ as soon as $\frac{n}{\log(4n)} \geq 168\, m$. In the sequel we take

$$n = \lfloor C_0 m \log(m+1) \rfloor,$$

with C_0 large enough, so that the previous inequality holds true. (One can take $C_0 = 10^4$, for instance.) Thanks to Lemma 16, we know that there exists a realization $x_1, \dots, x_n \in D_0$ of the random sampling such that the corresponding family $y_1, \dots, y_n$ satisfies

$$\left\| \frac{1}{n} \sum_{i=1}^{n} y_i y_i^* - E \right\|_{2\to 2} \leq \frac{1}{2}. \tag{8}$$

We fix such a sequence for the rest of the proof of Theorem 1.

4. Subsampling of infinite vectors

We now want to apply the solution to the Kadison-Singer problem, or specifically to Weaver's conjecture, to the sum of rank-one matrices

$$\frac{1}{n} \sum_{i=1}^{n} y_i y_i^*,$$

in order to find a subsampling of order m preserving the spectral properties of the sum. The original result comes from the celebrated paper [34] by Marcus, Spielman and Srivastava, and has already been applied numerous times in approximation theory, see for instance [7, 21, 27, 28, 37, 38, 40, 50]. However, the original subsampling strategy only works for finite matrices. The main result of this section is the following infinite-dimensional variant, that might be of independent interest.

Proposition 17. *There are absolute constants $c_1 \leq 43200$, $c_2 \geq 50$, $c_3 \leq 21600$, with the following properties. Let $n, m \in \mathbb{N}$ and $y_1, \dots, y_n$ be vectors from $\ell_2(\mathbb{N}_0)$ satisfying $\|y_i\|_2^2 \leq 2m$ and*

$$\left\| \frac{1}{n} \sum_{i=1}^{n} y_i y_i^* - \begin{pmatrix} I_m & 0 \\ 0 & \Lambda \end{pmatrix} \right\|_{2\to 2} \leq \frac{1}{2}, \tag{9}$$

for some Hermitian matrix Λ *with* $\|\Lambda\|_{2\to 2} \leq 1$*, where* $I_m \in \mathbb{C}^{m\times m}$ *denotes the identity.*
Then, there is a subset $J \subset \{1,\dots,n\}$ *with* $|J| \leq c_1 m$*, such that*

$$\left(\frac{1}{m}\sum_{i\in J} y_i y_i^*\right)_{<m} \geq c_2\, I_m \qquad \text{and} \qquad \frac{1}{m}\sum_{i\in J} y_i y_i^* \leq c_3\, I,$$

where $A_{<m} := (A_{k,l})_{k,l<m}$ *and* $A \leq B$ *denotes the Loewner order of Hermitian matrices* A *and* B*.*

The conclusion can be understood as an upper bound on the largest eigenvalue of $A = \sum_{i\in J} y_i y_i^*$ and a lower bound on the smallest eigenvalue of $A_{<m}$. Note that the constants in Proposition 17, and hence also the final sampling size, are independent of n, the original sampling size. The rest of this section is devoted to the proof of this proposition.

4.1. **Reduction to finite dimension.** Let U_0 be a matrix whose columns form an orthonormal basis of

$$\operatorname{span}\{(y_i)_{\geq m} \colon i = 1,\dots,n\} \subset \ell_2,$$

where $(y_i)_{\geq m} = ((y_i)_k)_{k\geq m}$. Clearly, U_0 has at most n columns. Then we have that $U_0^* U_0$ is the identity matrix and in particular the spectral norm of U_0 and U_0^* equals one. We set

$$U = \begin{pmatrix} I_m & 0 \\ 0 & U_0 \end{pmatrix},$$

which is a matrix that satisfies $U^*U = I_p$, where $p \leq m+n$, and therefore also U and U^* have unit norm. We choose vectors $z_i \in \mathbb{C}^p$ that satisfy $U z_i = y_i$ for all $i \leq n$. Such vectors exist since y_i is contained in the span of the columns of U. Then we also have $z_i = U^* U z_i = U^* y_i$.

Let $E = \begin{pmatrix} I_m & 0 \\ 0 & \Lambda \end{pmatrix}$ be the matrix from Proposition 17. We define

$$\hat{E} = U^*\, E\, U = \begin{pmatrix} I_m & 0 \\ 0 & E' \end{pmatrix} \qquad \text{where} \qquad \|E'\|_{2\to 2} \leq \|E\|_{2\to 2} \leq 1.$$

With the norm bounds on U and U^*, equation (9) gives

$$\left\|\frac{1}{n}\sum_{i=1}^n z_i z_i^* - \hat{E}\right\|_{2\to 2} = \left\|U^*\left(\frac{1}{n}\sum_{i=1}^n y_i y_i^* - E\right)U\right\|_{2\to 2} \leq \frac{1}{2}.$$

4.2. **Approximating the identity.** In addition to finite dimension, the result from [34] requires the matrix $\frac{1}{n}\sum_{i=1}^n z_i z_i^*$ to be close to the identity in spectral norm, and this is not ensured here. To mitigate this defect, we artificially add rank-one matrices $z_i z_i^* \in \mathbb{C}^{p\times p}$ for $i = n+1,\dots,q$ in the following way.

As $I_p - \hat{E}$ is positive semi-definite, we can decompose it as a sum of rank-one matrices

$$I_p - \hat{E} = \begin{pmatrix} 0 & 0 \\ 0 & I_{p-m} - E' \end{pmatrix} = \sum_{j=1}^{p-m} t_j t_j^*,$$

where $t_j \in \mathbb{C}^p$. We now choose

$$z_i = \sqrt{\frac{n}{n_{j(i)}}}\, t_{j(i)}, \quad n_j = \left\lceil \frac{n}{2m} \|t_j\|_2^2 \right\rceil,$$

with $j(i) \in \{1, \dots, p-m\}$ such that $\{z_i,\, i = n+1, \dots, q\}$ contains exactly n_j copies of each $\sqrt{n/n_j}\, t_j$. In this way, for $i > n$, the first m entries of z_i are zero since this is true for the t_j,

$$\|z_i\|_2^2 \le \frac{n}{n_{j(i)}} \|t_{j(i)}\|_2^2 \le 2m,$$

and

$$\begin{aligned} \left\| \frac{1}{n} \sum_{i=1}^{q} z_i z_i^* - I_p \right\|_{2\to 2} &= \left\| \frac{1}{n} \sum_{i=1}^{n} z_i z_i^* + \sum_{j=1}^{p-m} t_j t_j^* - I_p \right\|_{2\to 2} \\ &= \left\| \frac{1}{n} \sum_{i=1}^{n} z_i z_i^* - \hat{E} \right\|_{2\to 2} \le \frac{1}{2}. \end{aligned}$$

Remark 18. As $\|t_j\|_2^2 \le \|I_p - \hat{E}\|_{2\to 2} \le 1$, we count

$$q = n + \sum_{j=1}^{p-m} n_j \le n + \sum_{j=1}^{p-m} \left(1 + \frac{n}{2m}\right) \le n + (p-m)\frac{n}{m} = \frac{np}{m}.$$

Conversely, taking traces in $\mathbb{C}^{p\times p}$, we find

$$\frac{p}{2} = \operatorname{Tr}\left(\frac{1}{2} I_p\right) \le \operatorname{Tr}\left(\frac{1}{n} \sum_{i=1}^{q} z_i z_i^*\right) = \frac{1}{n} \sum_{i=1}^{q} \|z_i\|_2^2 \le \frac{2mq}{n}.$$

So, we obtain $n/m \ge q/p \ge n/4m$. Recall that, given m the dimension of the approximation space V_m, we took $n = \mathcal{O}(m \log m)$ initial sample points, and vectors z_i of size $p = \mathcal{O}(m \log m)$. Hence, the number of such vectors is $q = \mathcal{O}(m \log^2 m)$. Surprisingly, we do not use estimates on p and q in the rest of the argument.

Remark 19. In fact, we did not need an exponential speed of convergence in the concentration inequality. The reduction of the sample size to $\mathcal{O}(m)$ points works for any initial set of sampling points satisfying (8). If the cardinality of the initial sample is $n = m\,\ell(m)$, where $\ell(m)$ is any positive function of m, we get $p = \mathcal{O}(m\,\ell(m))$ and $q = \mathcal{O}(m\,\ell(m)^2)$.

4.3. **Reduction of the sample size.** We can now use the Kadison-Singer solution from [34] in an iterated way, as proposed in Lemma 3 of [40], and later used in [7, 27, 28, 32, 37, 38, 50]. The following lemma is obtained from Corollary B and Lemma 1 in [40].

Lemma 20. *Let $z_1, \dots, z_q \in \mathbb{C}^p$ with $\|z_i\|_2^2 \le \delta$ and*

$$\alpha I_p \le \sum_{i=1}^{q} z_i z_i^* \le \beta I_p$$

for some $\beta \ge \alpha > 100\,\delta > 0$. Then there is a partition of $\{1, \dots, q\}$ into sets $J_1, \dots, J_t$ such that, for all $s \le t$, we have

$$25\,\delta I_p \;\le\; \sum_{i \in J_s} z_i z_i^* \;\le\; 3600\,\frac{\beta}{\alpha}\,\delta I_p.$$

Proof. Since the matrix $M = \sum_{i=1}^{q} z_i z_i^*$ is positive, we may define $\tilde{z}_i = M^{-1/2} z_i$. Then we have $\sum_{i=1}^{q} \tilde{z}_i \tilde{z}_i^* = I_p$ and $\|\tilde{z}_i\|_2^2 \le \delta/\alpha =: \delta' < 1/100$. By Corollary B and Lemma 1 in [40], noting that the constant C from Lemma 1 is at most 36, we get a partition of $\{1, \dots, q\}$ into sets $J_1, \dots, J_t$ such that, for all $s \le t$, we have

$$25\,\delta'\, I_p \;\le\; \sum_{i \in J_s} \tilde{z}_i \tilde{z}_i^* \;\le\; 3600\,\delta'\, I_p.$$

Now, using

$$\sum_{i \in J_s} z_i z_i^* = M^{1/2} \sum_{i \in J_s} \tilde{z}_i \tilde{z}_i^*\, M^{1/2},$$

we get the statement. □

Note that one could obtain better constants by adapting the proof of Theorem 2.3 from [38]. In our case, we have $\delta = 2m$, $\alpha = n/2$ and $\beta = 3n/2$. The relation $\alpha > 100\,\delta$ is satisfied. We thus obtain

$$50\,m I_p \;\le\; \sum_{i \in J_s} z_i z_i^* \;\le\; 21600\,m I_p.$$

for every J_s from the partition. Moreover, the inequality

$$\frac{n}{2} I_p \;\le\; \sum_{i=1}^{q} z_i z_i^* \;=\; \sum_{s=1}^{t} \sum_{i \in J_s} z_i z_i^* \;\le\; 21600\,t m I_p$$

implies that one of the sets $J' = J_s$ from the partition must satisfy

$$|J' \cap \{1, \dots, n\}| \;\le\; \frac{n}{t} \;\le\; 43200\,m.$$

After applying Lemma 20 and removing the indices from $J' \cap \{n+1, \dots, q\}$ corresponding to artificially added vectors, we are left with a set $J := J' \cap \{1, \dots, n\}$ of cardinality

$$|J| \;\le\; 43200\,m.$$

It remains to show that the artificial vectors do not interfere with our desired properties. For this, recall that $(z_i)_k = (y_i)_k$ for $k < m$ and $i \le n$, whereas the first m entries of $z_i \in \mathbb{C}^p$ are zero for $i > n$. Hence,

$$\left(\sum_{i\in J} y_i y_i^*\right)_{<m} = \left(\sum_{i\in J'} z_i z_i^*\right)_{<m} \ge 50\, m\, I_m,$$

where we use a simple linear algebra fact on self-adjoint matrices A:

$$\lambda_{\min}(A_{<m}) = \inf_{\substack{z\in\mathbb{C}^p,\, \|z\|_2=1 \\ z_k=0 \text{ for } k\ge m}} z^* A z \ge \inf_{z\in\mathbb{C}^p,\, \|z\|_2=1} z^* A z = \lambda_{\min}(A).$$

Similarly, and using positive definiteness, we have

$$\sum_{i\in J} z_i z_i^* \le \sum_{i\in J'} z_i z_i^* \le 21600\, m\, I_p.$$

With the orthogonal transformation U from Section 4.1, we get

$$\Big\| \sum_{i\in J} y_i y_i^* \Big\|_{2\to 2} = \Big\| U \left(\sum_{i\in J} z_i z_i^*\right) U^* \Big\|_{2\to 2} \le \Big\| \sum_{i\in J} z_i z_i^* \Big\|_{2\to 2} \le 21600\, m.$$

This proves Proposition 17.

□

Remark 21. It would be an interesting improvement to use the result of Batson, Spielman and Srivastava, see [3], instead of [34] for the subsampling. This earlier paper is applied to approximation theory in e.g. [32, 39, 51] and more recently in [4]. It presents a slightly less powerful method, requiring additional weights, but comes with an almost linear algorithmic complexity, see [31], and much smaller constants, which could make the bound presented here sharp also in terms of numerical values.

Remark 22. We recently learned that it might be possible to use results from [12], which work directly in an infinite-dimensional setting, to save the reduction to a finite dimension in Section 4.1. However, as the core of our method is [34], we decided to keep our more direct deduction.

5. Proof of the main theorem

We now have all the tools for proving Theorem 1.

To obtain our sampling points, we combine (8) for our initial vectors $y_i \in \ell_2(\mathbb{I})$ with Proposition 17. Clearly, Proposition 17 stays true if we replace $\mathbb{N}_0$ by the possibly finite index set $\mathbb{I}$. We obtain points $x_1, \dots, x_n \in D_0$ with $n \le 43200\, m$ such that the vectors

$$(y_i)_k = \begin{cases} \rho_m(x_i)^{-1/2}\, b_k(x_i) & \text{if } 0 \le k < m, \\ \rho_m(x_i)^{-1/2}\, \gamma_m^{-1}\, \sigma_k\, b_k(x_i) & \text{if } m \le k < M, \end{cases}$$

satisfy

$$\left(\sum_{i=1}^{n} y_i y_i^*\right)_{<m} \geq 50\, m\, I,$$

and

$$\left(\sum_{i=1}^{n} y_i y_i^*\right)_{\geq m} \leq 21600\, m\, I,$$

where we use the notation $A_{\geq m} = (A_{k,l})_{k,l\geq m}$ for a matrix A.

As in earlier papers, we use the *weighted least squares estimator*

$$A_n(f) := \operatorname*{argmin}_{g\in V_m} \sum_{i=1}^{n} \frac{|g(x_i) - f(x_i)|^2}{\rho_m(x_i)}$$

with V_m and ρ_m as defined in Sections 2 and 3, respectively, see [27]. This algorithm may be written as

$$A_n(f) = \sum_{k=1}^{m} (G^+ N f)_k\, b_k$$

where $N\colon F \to \mathbb{C}^n$ with $N(f) := \big(\rho_m(x_i)^{-1/2} f(x_i)\big)_{i\leq n}$ is the *information mapping* and $G^+ \in \mathbb{C}^{m\times n}$ is the Moore-Penrose inverse of the matrix

$$G := \big(\rho_m(x_i)^{-1/2}\, b_k(x_i)\big)_{i\leq n,\, k\leq m} \in \mathbb{C}^{n\times m}.$$

Since we have the identity $\overline{G^*G} = (\sum_{i=1}^n y_i y_i^*)_{<m}$, the matrix G has full rank and the spectral norm of G^+ is bounded by $(50m)^{-1/2}$. In particular, the argmin in the definition of A_n is uniquely defined and A_n satisfies $A_n(f) = f$ for all $f \in V_m$.
Denoting with Q_m the L_2-orthogonal projection onto $\operatorname{span}\{b_k\colon k \geq m\}$, we obtain for any $f \in H$ that

$$\begin{aligned}
\|f - A_n(f)\|_{L_2}^2 &= \|f - P_m f\|_{L_2}^2 + \|P_m f - A_n(f)\|_{L_2}^2 \\
&= \|Q_m f\|_{L_2}^2 + \|A_n(f - P_m f)\|_{L_2}^2 \\
&= \|Q_m f\|_{L_2}^2 + \big\|G^+ N(f - P_m f)\big\|_{\ell_2^n}^2 \\
&\leq \sigma_m^2\, \|Q_m f\|_H^2 + \big\|G^+\big\|_{2\to 2}^2 \cdot \|N(f - P_m f)\|_{\ell_2^m}^2 .
\end{aligned}$$

By (7) we have $N(f - P_m f) = \Phi \xi_f$, where

$$\Phi = \big(\rho_m(x_i)^{-1/2} \sigma_k\, b_k(x_i)\big)_{i\leq n,\, k\geq m} \quad \text{and} \quad \xi_f = (\langle f, \sigma_k\, b_k\rangle_H)_{k\geq m}.$$

The matrix Φ satisfies

$$\overline{\Phi^*\Phi} = \gamma_m^2 \left(\sum_{i=1}^{n} y_i y_i^*\right)_{\geq m}$$

and therefore its spectral norm is bounded by $(21600\, m\, \gamma_m^2)^{1/2}$. Thus,

$$\|N(f - P_m f)\|_{\ell_2^m}^2 \leq 21600\, m\, \gamma_m^2 \|\xi_f\|_2^2 = 21600\, m\, \gamma_m^2\, \|Q_m f\|_H^2.$$

In summary, we obtain for all $1 \leq m < M$ the bound

$$\|f - A_n(f)\|_{L_2}^2 \leq 433 \max\left\{\sigma_m^2, \frac{1}{m}\sum_{k\geq m}\sigma_k^2\right\} \|Q_m f\|_H^2 . \tag{10}$$

for all $f \in H$ and some $n \leq 43200\, m$. Taking the supremum over $f \in F$ and using that

$$\max\left\{\sigma_m^2, \frac{1}{m}\sum_{k\geq m}\sigma_k^2\right\} \leq \frac{2}{m}\sum_{k\geq \lceil m/2\rceil}\sigma_k^2,$$

we obtain

$$g_{43200\, m}(F)^2 \leq \frac{866}{m}\sum_{k\geq \lceil m/2\rceil}\sigma_k^2.$$

This finishes the proof of Theorem 1 with $c = 43200 \cdot 866$.

□

In fact, equation (10) provides a local upper bound which is sometimes superior to Theorem 1. We therefore state it separately.

Theorem 23. *Let μ be a measure on a set D and let $F \subset L_2(\mu)$ be the unit ball of a separable RKHS H such that the finite trace assumption (1) holds. For $m \in \mathbb{N}$, let P_m be the orthogonal projection onto the span V_m of the singular vectors corresponding to the m largest singular values of the embedding of H into L_2. Then there exist $x_1, \ldots, x_n \in D$ and $\varphi_1, \ldots, \varphi_n \in V_m$, where $n \leq 43200m$, such that, for all $f \in H$,*

$$\Big\|f - \sum_{i=1}^{n} f(x_i)\,\varphi_i\Big\|_{L_2}^2 \leq 433 \max\left\{d_m(F)^2, \frac{1}{m}\sum_{k\geq m} d_k(F)^2\right\}\big\|f - P_m f\big\|_H^2.$$

Remark 24. For the purpose of Theorem 1 it was enough to bound $\|f - P_m f\|_H \leq \|f\|_H$. However, Theorem 23 will be of advantage later for the study of general classes since it is able to see additional decay of the Fourier coefficients $\langle f, b_k\rangle_{L_2}$ compared to the decay implied by $f \in H$. Note that faster decay of the Fourier coefficients often corresponds to higher smoothness of the function. In a certain sense, this means that the algorithm is universal. The error has the optimal rate of decay for any smoothness higher than the smoothness of H.

Remark 25. The condition on the point sets can also be given by finite matrices that are related to the kernel K of the Hilbert space. For this, let us define $K_m(x,y) := \sum_{k<m} b_k(x)b_k(y)$, and $R_m(x,y) := \sum_{k\geq m}\sigma_k^2\, b_k(x)b_k(y)$. The non-zero singular values of GG^* are the same as those of G^*G, and the non-zero singular values of $\Phi\Phi^*$ are the same as those of $\Phi^*\Phi$, where G and Φ are from above. Hence, the algorithm

A_n based on points $x_1, \dots, x_n$ satisfies the error bound above (up to a constant) if

$$c\,m \;\le\; \lambda_m\,(GG^*) \;=\; \lambda_m\left(\left(\frac{K_m(x_i,x_j)}{\sqrt{\rho_m(x_i)\rho_m(x_j)}}\right)_{i,j=1}^{n}\right)$$

and

$$\left(\frac{R_m(x_i,x_j)}{\sqrt{\rho_m(x_i)\rho_m(x_j)}}\right)_{i,j=1}^{n} \;=\; \Phi\Phi^* \;\le\; C\,m\,\gamma_m^2\,I$$

for some constants $c, C > 0$, where λ_m denotes the mth eigenvalue. It would be interesting to find a property that only involves the kernel K directly (instead of the truncated kernels above), or to verify that a similar property characterizes *good* point sets, in a way similar to Proposition 1 of [16] for integration.

5.1. **Proof of Corollary 2.** For the given bounds on the sampling numbers for sequences of polynomial decay, we only need to note that

$$\frac{1}{n}\sum_{k\ge n} k^{-a}\log^{-b} k \;\lesssim\; \begin{cases} n^{-a}\log^{-b} n & \text{if } a>1,\ b\in\mathbb{R},\\ n^{-a}\log^{-b+1} n & \text{if } a=1,\ b>1. \end{cases}$$

Hence, Corollary 2 immediately follows from Theorem 1, and the existence of F where the bounds are attained comes from (2), see [17]. □

6. General function classes

We now prove all results related to general function classes.

6.1. **Proof of Theorem 3.** We will make use of the following observation from [28, Lemma 3]. We copy its proof for completeness.

Lemma 26. *Let $F \subset L_2$ and let L_2 be infinite-dimensional. There is an orthonormal system $(b_k)_{k\in\mathbb{N}_0}$ in L_2 such that for all $m \ge 1$, the orthogonal projection P_m onto $V_m = \operatorname{span}\{b_k : k < m\}$ satisfies*

$$\sup_{f\in F}\|f - P_m f\|_{L_2} \;\le\; 2\,d_{\lfloor m/4\rfloor}(F). \tag{11}$$

Proof. Clearly it is enough to find an increasing sequence of subspaces of L_2,

$$U_1 \subseteq U_2 \subseteq U_3 \subseteq \dots, \qquad \dim(U_m) \le m,$$

such that the projection P_m onto U_m satisfies (11). By the definition of $d_k(F)$, $k \in \mathbb{N}_0$, there is a subspace $W_k \subset L_2$ of dimension k and a mapping $T_k\colon F \to W_k$ such that

$$\sup_{f\in F}\|f - T_k f\|_{L_2} \;\le\; 2\,d_k(F).$$

This is also true if $d_k(F) = 0$. We let U_m be the space that is spanned by the union of the spaces W_{2^ℓ} over all $\ell \in \mathbb{N}_0$ such that $2^\ell \leq m/2$. Note that U_m contains a subspace W_k with $k \geq \lfloor m/4 \rfloor$. Therefore, $P_m f$ is at least as close to f as $T_k f$ for some $k \geq \lfloor m/4 \rfloor$, which implies (11). □

We now turn to the proof of Theorem 3. The basic idea is to construct a suitable reproducing kernel Hilbert space H that contains a dense subset of F and apply Theorem 23 to this Hilbert space. It will be important to use the local bound from Theorem 23 instead of the global bound from Theorem 1.

Proof of Theorem 3. Without loss of generality, we assume that L_2 is infinite-dimensional. Moreover, we assume that $d_k(F)$ is finite for $k \geq k_0$ and that $(d_k(F))_{k \geq k_0} \in \ell_p$. Otherwise, the statement is trivial.

By Lemma 26, there is an orthonormal system $(b_k)_{k \in \mathbb{N}_0}$ such that (11) is satisfied for all $m \in \mathbb{N}$. We will consider b_k as a function, where we fix an arbitrary representer from the equivalence class in L_2. We call

$$\hat{f}(k) := \langle f, b_k \rangle_{L_2}$$

the kth Fourier coefficient of f. Moreover, we fix a countable dense subset F_0 of F and set $\sigma_k = \max\{1, k\}^{-\alpha}$ for all $k \in \mathbb{N}_0$ and some $\alpha \in (1/2, 1/p)$. Then we have $(\sigma_k) \in \ell_2$.

We now want to define a RKHS on a set $D_0 \subset D$, with $\mu(D \setminus D_0) = 0$, which admits the orthonormal basis $(\sigma_k b_k)$ and contains the set F_0. Such a Hilbert space will have the reproducing kernel

$$K(x, y) \;=\; \sum_{k \in \mathbb{N}_0} \sigma_k^2 b_k(x) \overline{b_k(y)}.$$

To find a suitable set D_0, we first note that

$$\int_D K(x, x) \, d\mu(x) \;=\; \sum_{k \in \mathbb{N}_0} \sigma_k^2 < \infty \tag{12}$$

and thus $K(x, x)$ is finite for all $x \in D \setminus E$ with a null set $E \subset D$. Moreover, for all $f \in F_0$, we have

$$\sum_{k \geq 1} k \, |\hat{f}(k)|^2 \;=\; \sum_{n \geq 0} \sum_{k > n} |\hat{f}(k)|^2 \;=\; \sum_{n \geq 0} \|f - P_n f\|_{L_2}^2 < \infty,$$

where we use (11) and the assumptions on F. The Rademacher-Menchov Theorem, see e.g. [45], now implies that the Fourier series of f at x converges to $f(x)$ for all $x \in D \setminus E_f$ with a null set $E_f \subset D$. We put $D_0 := D \setminus E_0$, where $E_0 := E \cup \bigcup_{f \in F_0} E_f$ is a null set. Then for all $x \in D_0$ and $f \in F_0$, we have

$$K(x, x) \;< \infty \quad \text{and} \quad f(x) \;=\; \sum_{k \in \mathbb{N}_0} \hat{f}(k) \, b_k(x).$$

We now define the space H as the set of all square-integrable functions $f\colon D_0 \to \mathbb{C}$ which are point-wise represented by their Fourier series $\sum_k \hat{f}(k)\, b_k$ and which satisfy

$$\|f\|_H^2 := \sum_{k\in\mathbb{N}_0} \frac{|\hat{f}(k)|^2}{\sigma_k^2} < \infty.$$

Then H is a separable reproducing kernel Hilbert space on D_0 since

$$|f(x)|^2 \le K(x,x)\|f\|_H^2 \quad \text{for all } x \in D_0 \text{ and } f \in H,$$

and $(\sigma_k b_k)_{k\in\mathbb{N}_0}$ is an orthonormal basis of H. The reproducing kernel is K, which has finite trace from (12).

We now show that F_0 (with functions restricted to D_0) is a subset of H. Recall that any $f \in F_0$ is point-wise represented by its Fourier series. Moreover, note that the Kolmogorov widths of F_0 and F are the same. We use

$$d_{2m}(F) = \big(d_{2m}(F)^p\big)^{1/p} \le \left(\frac{1}{m}\sum_{k\ge m} d_k(F)^p\right)^{1/p}$$

and obtain for any $m \in 8\mathbb{N}$ and $f \in F_0$ that

$$\begin{aligned}
\|f - P_m f\|_H^2 &= \sum_{k\ge m} k^{2\alpha}\,|\hat{f}(k)|^2 \le \sum_{\ell\in\mathbb{N}_0} \big(m2^{\ell+1}\big)^{2\alpha} \sum_{k=m2^\ell}^{m2^{\ell+1}-1} |\hat{f}(k)|^2 \\
&\le 4\sum_{\ell\in\mathbb{N}_0} \big(m2^{\ell+1}\big)^{2\alpha}\, d_{m2^{\ell-2}}(F)^2 \\
&\le 4\sum_{\ell\in\mathbb{N}_0} \big(m2^{\ell+1}\big)^{2\alpha} \left(\frac{1}{m2^{\ell-3}} \sum_{k\ge m2^{\ell-3}} d_k(F)^p\right)^{2/p} \\
&\le\ 2^{2+2\alpha+6/p} m^{2\alpha-2/p} \sum_{\ell\in\mathbb{N}_0} 2^{(2\alpha-2/p)\ell} \left(\sum_{k\ge m/8} d_k(F)^p\right)^{2/p}.
\end{aligned}$$

The last expression is finite for $m \ge 8k_0$, since $2\alpha - 2/p < 0$. This implies that $f \in H$ and

$$\|f - P_m f\|_H \ \le\ C\, m^\alpha \left(\frac{1}{m}\sum_{k\ge m/8} d_k(F)^p\right)^{1/p}, \tag{13}$$

where $C > 0$ only depends on $p \in (0,2)$ and $\alpha \in (\frac{1}{2}, \frac{1}{p})$.

We now apply Theorem 23 to the newly constructed Hilbert space H to find $n \le 43200\, m$ and a linear algorithm A_n of the form

$$A_n(f) \ =\ \sum_{i=1}^n f(x_i) g_i, \quad x_i \in D_0,\ g_i \in L_2,$$

such that

$$\|f - A_n f\|_{L_2(D_0,\mu)}^2 \le 433 \max\left\{\sigma_m^2, \frac{1}{m}\sum_{k\ge m}\sigma_k^2\right\}\|f - P_m f\|_H^2 \tag{14}$$

for all $f \in H$ and thus, for all $f \in F_0$. Clearly, in the last inequality, D_0 can be replaced with D. If we now insert the estimate (13) and the estimate

$$\max\left\{\sigma_m^2, \frac{1}{m}\sum_{k\ge m}\sigma_k^2\right\} \lesssim m^{-2\alpha}, \tag{15}$$

into (14), we obtain that

$$\|f - A_n f\|_{L_2}^2 \le \left(\frac{\tilde{c}_p}{m}\sum_{k\ge m/8} d_k(F)^p\right)^{2/p}$$

for all $f \in F_0$ and some $\tilde{c}_p > 0$ that only depends on p. Since F_0 is dense in F and both $\mathrm{id}\colon F \to L_2$ and $A_n\colon F \to L_2$ are continuous, the last bound is true for all $f \in F$. This finishes the proof of Theorem 3 with $c_p = 43200\max(\tilde{c}_p, 8)$.

□

6.2. **The boundary case.** We provide a variant of Theorem 3 under a weaker condition than $(d_k(F)) \in \ell_p$ for $p < 2$. In fact, we show that the condition $\big((\log k)^s d_k(F)\big) \in \ell_2$ for some $s > 1/2$ is enough for a comparison of the sampling and the Kolmogorov widths, while the same assumption for $s = 1/2$ is not enough, see Example 30.

Theorem 27. *Let $s > 1/2$. There is a universal constant $c \in \mathbb{N}$ and a constant $c_s > 0$, depending only on s, such that for every F and μ that satisfy Assumption A and all $m \ge 2$,*

$$g_{cm}(F)^2 \le c_s\, m^{-1}\log^{-2s+1} m \sum_{k\ge m} d_k(F)^2 \cdot \log^{2s} k.$$

Proof. The proof follows the same lines as the proof of Theorem 3. The only difference is that we now choose $\sigma_k = k^{-1/2}\log^{-s} k$ for $k \ge 2$. Then, inequality (13) becomes

$$\begin{aligned}\|f - P_m f\|_H^2 &= \sum_{k\ge m}|\hat{f}(k)|^2 k\log^{2s}(k) \le \sum_{k\ge m}|\hat{f}(k)|^2 \sum_{m\le r\le 2k}\log^{2s}(r)\\ &\le \sum_{r\ge m}\log^{2s}(r)\sum_{k\ge r/2}|\hat{f}(k)|^2 \le 4\sum_{r\ge m}\log^{2s}(r)\, d_{\lfloor r/8\rfloor}(F)^2\\ &\le 32\sum_{k\ge\lfloor m/8\rfloor}\log^{2s}(8k+7)\, d_k(F)^2.\end{aligned}$$

Likewise, inequality (15) becomes

$$\max\left\{\sigma_m^2,\, \frac{1}{m}\sum_{k\geq m}\sigma_k^2\right\} \lesssim m^{-1}\log^{-2s+1} m$$

and the stated inequality is obtained. □

6.3. **Proof of Corollary 4.** Using the same bound as in the proof of Corollary 2, the case $\alpha > 1/2$ immediately follows from Theorem 3 if we choose $1/\alpha < p < 2$, and the case $\alpha = 1/2$, $\beta > 1$ from Theorem 27 if we choose $1/2 < s < \beta - 1/2$.

All bounds are attained with the same classes F as in Corollary 2 for the first case, and with the constructions from the next section for the two other cases.

□

7. Examples

We first apply Theorem 1 to tensor product spaces.

Example 28. Let H be a RKHS on D that is compactly embedded into L_2 and let F be its unit ball. We consider L_2-approximation on the unit ball F_d of the d-fold tensor product H_d of H, which is a RKHS on the domain D^d. We assume that $g_n(F) \lesssim n^{-\alpha}$ for some $\alpha > 0$. The famous Smolyak algorithm, see [47], gives the estimate

$$g_n(F_d) \lesssim n^{-\alpha}\log^{(\alpha+1)(d-1)} n. \tag{16}$$

An example of such tensor product spaces are the spaces of dominating mixed smoothness $\alpha > 1/2$, see [11]. For these spaces, it is known that the error bound (16) for the Smolyak algorithm can be improved [46]; the exponent of the logarithm can be reduced to $(\alpha+1/2)(d-1)$. With Corollary 2 and known results on the approximation numbers of tensor product operators, see [2, 36], we now obtain

$$g_n(F_d) \lesssim n^{-\alpha}\log^{\alpha(d-1)} n \quad \text{if} \quad \alpha > 1/2. \tag{17}$$

This bound is asymptotically optimal for the spaces of mixed smoothness, see [48, Theorem 1] or [49, Theorem 6.4.3]. More generally, it is known that $d_n(F) \asymp n^{-\alpha}$ implies $d_n(F_d) \asymp n^{-\alpha}\log^{\alpha(d-1)} n$ (see e.g. [22]) and therefore the asymptotic bound (17) is optimal whenever the approximation numbers in the univariate case are of order $n^{-\alpha}$. Let us note, however, that also preasymptotic estimates on the sampling numbers (say, for $n < d^d$) are of interest, especially if the dimension d is high, see [22, 29, 58].

□

We now present two examples that show that our upper bounds cannot be improved without further assumptions on the class F.

First, we show that the worst possible behavior of the sampling numbers in the case $d_n(F) \lesssim n^{-1/2}\log^{-\beta} n$ with $\beta > 1$ is indeed $n^{-1/2}\log^{-\beta+1} n$.

Example 29. For $\ell \in \mathbb{N}_0$ and $k \in \{1, \dots, 2^\ell\}$ define the interval $I_{\ell,k} = [(k-1)2^{-\ell}, k2^{-\ell})$ and denote with $\chi_{\ell,k}$ the indicator function of $I_{\ell,k}$. Let $\beta > 1$. We set

$$\mathcal{C}_\beta := \left\{ \mathbf{c} := (c_{\ell,k})_{\ell\in\mathbb{N}_0,\, 1\le k\le 2^\ell} \;\middle|\; \sum_{k=1}^{2^\ell} |c_{\ell,k}|^2 \le (\ell+1)^{-2\beta} \text{ for all } \ell \in \mathbb{N}_0 \right\}$$

and consider the class

$$F_\beta := \left\{ f_{\mathbf{c}} := \sum_{\ell\in\mathbb{N}_0} \sum_{k=1}^{2^\ell} c_{\ell,k}\chi_{\ell,k} \;\middle|\; \mathbf{c} \in \mathcal{C}_\beta \right\}.$$

Note that the series $f_{\mathbf{c}}$ converge uniformly, since the inner sum is bounded by $(\ell+1)^{-\beta}$. If F_β is equipped with the maximum distance on $[0,1)$, it is a separable metric space, function evaluation is continuous, and the embedding in $L_2([0,1))$ is continuous.

For every $L \in \mathbb{N}_0$, the span V_L of the functions $\chi_{\ell,k}$ with $\ell \le L$ has dimension 2^L. If P_L is the L_2-orthogonal projection onto V_L, we have for all $\mathbf{c} \in \mathcal{C}_\beta$ that

$$\begin{aligned}\left\|f_{\mathbf{c}} - P_L f_{\mathbf{c}}\right\|_2 &\le \left\| \sum_{(\ell,k)\colon \ell>L} c_{\ell,k}\chi_{\ell,k} \right\|_2 \le \sum_{\ell>L} \left\| \sum_{k=1}^{2^\ell} c_{\ell,k}\chi_{\ell,k} \right\|_2 \\ &= \sum_{\ell>L} \left(\sum_{k=1}^{2^\ell} c_{\ell,k}^2 \|\chi_{\ell,k}\|_2^2 \right)^{1/2} \le \sum_{\ell>L} 2^{-\ell/2}(\ell+1)^{-\beta} \lesssim 2^{-L/2}L^{-\beta},\end{aligned}$$

and thus

$$d_{2^L}(F_\beta) \lesssim 2^{-L/2}L^{-\beta},$$

or equivalently

$$d_n(F_\beta) \lesssim n^{-1/2}\log^{-\beta} n.$$

We now show a lower bound for the sampling numbers. Let $x_1, \dots, x_n \in [0,1)$. For all $\ell \in \mathbb{N}_0$, we let J_ℓ be the set of indices $1 \le k \le 2^\ell$ such that $I_{\ell,k}$ contains at least one of these points. Clearly, the cardinality of J_ℓ is at most n. We choose $L \in \mathbb{N}_0$ of order $\log n$ and define

$$f_L := \sum_{\ell>L} |J_\ell|^{-1/2}(\ell+1)^{-\beta} \sum_{k\in J_\ell} \chi_{\ell,k}.$$

This function is contained in F_β and for all $i \le n$, we have

$$h := f_L(x_i) = \sum_{\ell>L} |J_\ell|^{-1/2}(\ell+1)^{-\beta} \gtrsim n^{-1/2}\log^{-\beta+1} n,$$

where h is independent of i. On the other hand, as shown by our previous calculation,

$$\left| \int_0^1 f_L(x)\,dx \right| \le \left\| f_L \right\|_2 \lesssim 2^{-L/2} L^{-\beta} \lesssim n^{-1/2} \log^{-\beta} n.$$

Thus, if we set $f = h - f_L$, the function is contained in F_β, vanishes at all points $x_1, \dots, x_n$, and satisfies

$$\|f\|_2 \ge \int_0^1 f(x)\,dx \ge h - \left| \int_0^1 f_L(x)\,dx \right| \gtrsim n^{-1/2} \log^{-\beta+1} n.$$

This shows $g_n(F_\beta) \gtrsim n^{-1/2} \log^{-\beta+1} n$.

□

The next example shows that, in the case $d_n(F) \lesssim n^{-1/2} \log^{-\beta} n$ with $\beta \le 1$, no general statement on the sampling numbers is possible.

Example 30. Similar to Example 29, we define

$$\mathcal{C} := \left\{ \mathbf{c} \;\middle|\; \sum_{k=1}^{2^\ell} |c_{\ell,k}|^2 \le (\ell+1)^{-2} \log(\ell+e)^{-2} \text{ for all } \ell \in \mathbb{N}_0 \right\}$$

and consider the class

$$F := \left\{ f_{\mathbf{c}} := \sum_{\ell \in \mathbb{N}_0} \sum_{k=1}^{2^\ell} c_{\ell,k} \chi_{\ell,k} \;\middle|\; \mathbf{c} \in \mathcal{C},\, \mathbf{c} \text{ finite} \right\}.$$

The finiteness of the sequences ensures that F, equipped with the maximum distance, is still a separable metric space, where function evaluation is continuous, and the embedding in $L_2([0,1))$ is continuous. As above, we obtain

$$d_n(F) \lesssim n^{-1/2} (\log n)^{-1} (\log\log n)^{-1}.$$

In particular, we have $(d_n(F) \log^{1/2} n) \in \ell_2$. On the other hand, given $x_1, \dots, x_n$ and $\varepsilon > 0$, we choose $L \in \mathbb{N}_0$ with

$$\sum_{\ell > L} 2^{-\ell/2} (\ell+1)^{-1} (\log(\ell+e))^{-1} \le \varepsilon,$$

define the sets J_ℓ as above, and choose $N \in \mathbb{N}_0$ such that

$$h := \sum_{\ell=L+1}^{N} |J_\ell|^{-1/2} (\ell+1)^{-1} (\log(\ell+e))^{-1} \ge 1.$$

The function

$$f_L := \frac{1}{h} \sum_{\ell=L+1}^{N} |J_\ell|^{-1/2} (\ell+1)^{-1} (\log(\ell+e))^{-1} \sum_{k \in J_\ell} \chi_{\ell,k},$$

is contained in F, its integral is at most ε, and it satisfies $f_L(x_i) = 1$ for all $i \leq n$. Then $f = 1 - f_L$ is contained in F, vanishes at all points $x_1, \dots, x_n$, and satisfies

$$\|f\|_2 \geq \int_0^1 f(x)\,dx \geq 1 - \left| \int_0^1 f_L(x)\,dx \right| \geq 1 - \varepsilon.$$

This shows $g_n(F) \geq 1$ for all $n \in \mathbb{N}_0$.

□

We note that the lower bounds in Example 29 and 30 already hold for the easier problem of numerical integration on F_β. Thus, the upper bounds from Corollary 4 are also sharp for the minimal error of quadrature rules on probability spaces.

References

[1] S. T. Ali, J.-P. Antoine, J.-P. Gazeau, *Coherent states, wavelets, and their generalizations*. Second edition. Theoretical and Mathematical Physics. Springer, New York, 2014.

[2] K. I. Babenko, *About the approximation of periodic functions of many variable trigonometric polynomials*. Doklady Akademii Nauk SSSR, 32, 247–250, 1960.

[3] J. Batson, D.A. Spielman, and N. Srivastava, *Twice-Ramanujan Sparsifiers*, SIAM Review, 56, 315-334, 2014.

[4] F. Bartel, M. Schäfer and T. Ullrich, *Constructive subsampling of finite frames with applications in optimal function recovery*, arXiv:2202.12625, 2022.

[5] M. Bownik, *Continuous frames and the Kadison-Singer problem*. Coherent states and their applications, 63-88. Springer, Cham, 2018.

[6] O. Christensen, *An introduction to frames and Riesz bases*, Applied and Numerical Harmonic Analysis, Birkhäuser Boston, Inc., Boston, MA, 2003.

[7] A. Cohen and M. Dolbeault, *Optimal pointwise sampling for L_2 approximation*, Journal of Complexity, 68, 101602, 2022.

[8] A. Cohen and G. Migliorati, *Optimal weighted least squares methods,* SMAI Journal of Computational Mathematics, 3, 181-203, 2017.

[9] F. Cucker and S. Smale, *On the mathematical foundations of learning*, Bulletin of the American mathematical society, 39(1), 1-49, 2002.

[10] R.A. DeVore and G.G. Lorentz, *Constructive approximation*, Vol. 303. Springer Science & Business Media, 1993.

[11] D. Dũng, V. N. Temlyakov and T. Ullrich. Hyperbolic Cross Approximation. *Advanced Courses in Mathematics - CRM Barcelona*. Springer International Publishing, 2018.

[12] D. Freeman and D. Speegle *The discretization problem for continuous frames*, Advances in Mathematics 345, 784-813, 2019.

[13] A. Hinrichs, E. Novak and J. Vybiral, *Linear information versus function evaluations for L_2-approximation*, Journal of approximation theory, 153(1), 97-107, 2008.

[14] A. Hinrichs, D. Krieg, E. Novak, J. Prochno and M. Ullrich, *Random sections of ellipsoids and the power of random information*, Transactions of the American Mathematical Society 374(12), 8691-8713, 2021.

[15] A. Hinrichs, D. Krieg, E. Novak, J. Prochno, and M. Ullrich. On the power of random information. In F. J. Hickernell and P. Kritzer, editors, *Multivariate Algorithms and Information-Based Complexity*, 43-64. De Gruyter, Berlin/Boston, 2020.

[16] A. Hinrichs, D. Krieg, E. Novak and J. Vybiral, *Lower bounds for the error of quadrature formulas for Hilbert spaces*, Journal of Complexity, 65, 101544, 2021, https://doi.org/10.1016/j.jco.2020.101544.

[17] A. Hinrichs, D. Krieg, E. Novak and J. Vybiral, *Lower bounds for integration and recovery in L_2*, Journal of Complexity, 101662, 2022, https://doi.org/10.1016/j.jco.2022.101662.

[18] A. Hinrichs, J. Prochno, and M. Sonnleitner, *Random sections of ℓ_p-ellipsoids, optimal recovery and Gelfand numbers of diagonal operators*, arXiv:2109.14504, 2021.

[19] L. Kaemmerer, T. Ullrich, and T. Volkmer, *Worst case recovery guarantees for least squares approximation using random samples*, Constructive Approximation, 54(2), 295-352, 2021.

[20] B. Kashin, E. Kosov, I. Limonova, and V. Temlyakov, *Sampling discretization and related problems*, Journal of Complexity, 101653, 2022, https://doi.org/10.1016/j.jco.2022.101653.

[21] D. Krieg, *Optimal Monte Carlo methods for L_2-approximation*, Constructive Approximation, 49, 385–403, 2019.

[22] D. Krieg, *Tensor power sequences and the approximation of tensor product operators*, Journal of Complexity, 44, 30–51, 2018.

[23] D. Krieg, E. Novak, and M. Sonnleitner, *Recovery of Sobolev functions restricted to iid sampling*, Mathematics of Computation, 91, 2715–2738, 2022.

[24] D. Krieg, P. Siedlecki, M. Ullrich, and H. Woźniakowski, *Exponential tractability of L_2-approximation with function values*, arXiv:2205.04141, 2022.

[25] D. Krieg and M. Sonnleitner, *Random points are optimal for the approximation of Sobolev functions*, arXiv:2009.11275, 2022.

[26] D. Krieg and M. Sonnleitner, *Function recovery on manifolds using scattered data*, arXiv:2109.04106, 2022.

[27] D. Krieg and M. Ullrich, *Function values are enough for L_2-approximation*, Foundations of Computational Mathematics, 21(4), 1141-1151, 2021.

[28] D. Krieg and M. Ullrich, *Function values are enough for L_2-approximation: Part II*, Journal of Complexity, 66, 101569, 2021, https://doi.org/10.1016/j.jco.2021.101569.

[29] T. Kühn, W. Sickel, and T. Ullrich, *Approximation of mixed order Sobolev functions on the d-torus – asymptotics, preasymptotics and d-dependence*. Constructive Approximation, 42, 353–398, 2015.

[30] F. Kuo, G. W. Wasilkowski, and H. Woźniakowski. *On the power of standard information for multivariate approximation in the worst case setting*, Journal of Approximation Theory, 158, 97–125, 2009.

[31] Y. T. Lee and H. Sun, *Constructing linear-sized spectral sparsification in almost-linear time*, SIAM Journal on Computing, 47(6), 2315-2336, 2018.

[32] I. Limonova and V. Temlyakov, *On sampling discretization in L_2*, Journal of Mathematical Analysis and Applications, 515(2), 126457, 2022.

[33] W. Lu and H. Wang, *On the power of standard information for tractability for L_2-approximation in the randomized setting*, Journal of Complexity, 70, 101618, 2022.

[34] A. Marcus, D. Spielman and N. Srivastava, *Interlacing families II: Mixed characteristic polynomials and the Kadison-Singer problem*, Annals of Mathematics, 327-350, 2015, http://dx.doi.org/10.4007/annals.2015.182.1.8.

[35] S. Mendelson and A. Pajor, *On singular values of matrices with independent rows*, Bernoulli, 12(5), 761-773, 2006.

[36] B. S. Mityagin, *Approximation of functions in L^p and C on the torus*. Mathematical Notes, 58, 397–414, 1962.

[37] M. Moeller and T. Ullrich, *L_2-norm sampling discretization and recovery of functions from RKHS with finite trace*, Sampling Theory, Signal Processing, and Data Analysis, 19(2), 1-31, 2021.

[38] N. Nagel, M. Schäfer, and T. Ullrich, *A new upper bound for sampling numbers*, Foundations of Computational Mathematics, 22(2), 445-468, 2022.

[39] S. Nitzan, A. Olevskii and A. Ulanovskii, *A few remarks on sampling of signals with small spectrum*, Proceedings of the Steklov Institute of Mathematics, 280(1), 240-247, 2013.

[40] S. Nitzan, A. Olevskii and A. Ulanovskii, *Exponential frames on unbounded sets*, Proceedings of the American Mathematical Society, vol. 144, no 1, 109-118, 2016, http://dx.doi.org/10.1090/proc/12868.

[41] E. Novak and H. Woźniakowski. *Tractability of multivariate problems. Volume I: Linear information*, volume 6 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2008.

[42] E. Novak and H. Woźniakowski. *Tractability of multivariate problems. Volume II: Standard information for functionals*, volume 12 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2010.

[43] E. Novak and H. Woźniakowski, *Tractability of multivariate problems. Volume III: Standard information for operators*, volume 18 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2012.

[44] E. Novak and H. Woźniakowski. *Tractability of multivariate problems for standard and linear information in the worst case setting: part I*, Journal of Approximation Theory, 207, 177–192, 2016.

[45] R. Salem. *A new proof of a theorem of Menchoff*, Duke Mathematical Journal, 8(2), 269–272, 1941.

[46] W. Sickel and T. Ullrich, *Smolyak's algorithm, sampling on sparse grids and function spaces of dominating mixed smoothness*, East Journal on Approximations, 13(4), 387–425, 2007.

[47] S. A. Smolyak, *Quadrature and interpolation formulas for tensor products of certain classes of functions*, Soviet Mathematics Doklady, 4, 240–243, 1963.

[48] V. N. Temlyakov. *On a way of obtaining lower estimates for the errors of quadrature formulas*, Mathematics of the USSR-Sbornik, 71(1), 247, 1992.

[49] V. N. Temlyakov. *Multivariate Approximation*, volume 32 of *Cambridge Monographs on Applied and Computational Mathematics*. Cambridge University Press, 2018.

[50] V. N. Temlyakov, *On optimal recovery in L_2*, Journal of Complexity, 65, 101545, 2020, https://doi.org/10.1016/j.jco.2020.101545.

[51] V. N. Temlyakov, *The Marcinkiewicz-type discretization theorems*, Constructive Approximation, 48, 337-369, 2018.

[52] V. N. Temlyakov, T. Ullrich. *Approximation of functions with small mixed smoothness in the uniform norm*, Journal of Approximation Theory, 277, 105718, 2022.

[53] V. N. Temlyakov and T. Ullrich, *Bounds on Kolmogorov widths and sampling recovery for classes with small mixed smoothness*, Journal of Complexity, 67, 101575, 2021.

[54] J. F. Traub, H. Woźniakowski. *A general theory of optimal algorithms*. Academic Press, 1980.

[55] M. Ullrich, *On the worst-case error of least squares algorithms for L_2-approximation with high probability*, Journal of Complexity, 60, 2020.
[56] J. Vybíral, *A variant of Schur's product theorem and its applications*, Advances in Mathematics, 368, 107140, 2020.
[57] M. J. Wainwright. *High-Dimensional statistics. A non-asymptotic viewpoint.* Cambridge University Press, 2019.
[58] G. W. Wasilkowski and H. Woźniakowski, *Explicit cost bounds of algorithms for multivariate tensor product problems*, Journal of Complexity, 11, 1–56, 1995.
[59] G. W. Wasilkowski and H. Woźniakowski. *On the power of standard information for weighted approximation*, Foundations of Computational Mathematics, 1(4), 417-434, 2001.
[60] G.W. Wasilkowski and H. Woźniakowski, *The power of standard information for multivariate approximation in the randomized setting*, Mathematics of Computation, 76, 965-988, 2007.
[61] H. Wendland, *Scattered data approximation*, Vol. 17, Cambridge university press, 2004.