



HAL
open science

Single mRNA detection of Wnt signaling pathway in the human limbus

Clémence Bonnet, Maxime Ruiz, Sheyla Gonzalez, Chi-Hong Tseng,
Jean-Louis Bourges, Francine Behar-Cohen, Sophie X Deng

► **To cite this version:**

Clémence Bonnet, Maxime Ruiz, Sheyla Gonzalez, Chi-Hong Tseng, Jean-Louis Bourges, et al.. Single mRNA detection of Wnt signaling pathway in the human limbus. *Experimental Eye Research*, 2023, 229, 10.1016/j.exer.2022.109337 . hal-04088567

HAL Id: hal-04088567

<https://u-paris.hal.science/hal-04088567>

Submitted on 4 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Research article

Single mRNA detection of Wnt signaling pathway in the human limbus

Clémence Bonnet^{a,b}, Maxime Ruiz^a, Sheyla Gonzalez^a, Chi-Hong Tseng^c, Jean-Louis Bourges^b, Francine Behar-Cohen^b, Sophie X. Deng^{a,d,*}^a Stein Eye Institute, Department of Ophthalmology, David Geffen School of Medicine, University of California, Los Angeles, CA, USA^b INSERM, UMRS1138, Team 17, From Physiopathology of Ocular Diseases to Clinical Development, Centre de Recherche des Cordeliers, Paris University, And Cornea Department, Cochin Hospital, AP-HP, F-75014, Paris, France^c David Geffen School of Medicine, Division of General Internal Medicine and Health Services Research, University of California, Los Angeles, USA^d Molecular Biology Institute, University of California, Los Angeles, CA, USA

ARTICLE INFO

Keywords:

Cornea
Limbus
frizzled7
Limbal stem cell
Limbal stem cell deficiency
RNA in situ hybridization
RNAscope ISH
Wnt signaling pathway

ABSTRACT

Limbal epithelial stem/progenitor cells (LSCs) are adult stem cells located at the limbus, tightly regulated by their close microenvironment. It has been shown that Wnt signaling pathway is crucial for LSCs regulation. Previous differential gene profiling studies confirmed the preferential expression of specific Wnt ligands (WNT2, WNT6, WNT11, WNT16) and Wnt inhibitors (DKK1, SFRP5, WIF1, FRZB) in the limbal region compared to the cornea. Among all frizzled receptors, frizzled7 (Fzd7) was found to be preferentially expressed in the basal limbal epithelium. However, the exact localization of Wnt signaling molecules-producing cells in the limbus remains unknown. The current study aims to evaluate the in situ spatial expression of these 4 Wnt ligands, 4 Wnt inhibitors, and Fzd7. Wnt ligands, DKK1, and Fzd7 expression were scattered within the limbal epithelium, at a higher abundance in the basal layer than the superficial layer. SFRP5 expression was diffuse among the limbal epithelium, whereas WIF1 and FRZB expression was clustered at the basal limbal epithelial layer corresponding to the areas of high levels of Fzd7 expression. Quantitation of the fluorescence intensity showed that all 4 Wnt ligands, 3 Wnt inhibitors (WIF1, DKK1, FRZB), and Fzd7 were highly expressed at the basal layer of the limbus, then in a decreasing gradient toward the superficial layer ($P < 0.05$). The expression levels of all 4 Wnt ligands, FRZB, and Fzd7 in the basal epithelial layer were higher in the limbus than the central cornea ($P < 0.05$). All 4 Wnt ligands, 4 Wnt inhibitors, and Fzd7 were also highly expressed in the limbal stroma immediately below the epithelium but not in the corneal stroma ($P < 0.05$). In addition, Fzd7 had a preferential expression in the superior limbus compared to other limbal quadrants ($P < 0.05$). Taken together, the unique expression patterns of the Wnt molecules in the limbus suggests the involvement of both paracrine and autocrine effects in LSCs regulation, and a fine balance between Wnt activators and inhibitors to govern LSC fate.

1. Introduction

The basal epithelium of the human limbus contains a distinct population of adult epithelial stem cells called limbal stem/progenitor cells (LSCs) (Davanger, 1971; Tseng, 1989). LSCs closely interact with their microenvironment, referred to as the stem cell niche, consisting of stromal cells, melanocytes, and extracellular matrix, in a highly vascularized and innervated stroma (Dziasko, 2016a; Scadden, 2006). These interactions confer dual functions on the limbus, i.e., formation of a barrier between the corneal and conjunctival epithelium, and renewal of

corneal epithelial cells (Cotsarelis, 1989). Both functions are necessary to maintain homeostasis of the corneal epithelium, the integrity of the ocular surface, and visual function under physiologic conditions. Dysfunction or deficiency of LSCs and/or their niche can lead to limbal stem cell deficiency (LSCD), a condition that can cause corneal blindness (Bonnet, 2021d; Deng, 2019). Therapies of LSCD include surgical management when the LSC dysfunction cannot be medically treated and/or when the LSC pool needs to be restored (Deng, 2020b). Surgical techniques have been optimized to reduce the size of the donor's tissue required (Basu, 2016), and cell-based therapies from ex vivo LSC

Abbreviations: DKK1, dickkopf WNT Signaling Pathway Inhibitor 1; FRZB, frizzled-related protein; FZD, frizzled; ISH, in-situ hybridization; LSC, limbal stem cell; LSCD, limbal stem cell deficiency; SFRP5, secreted frizzled-related protein 5; WIF1, WNT inhibitory factor 1.

* Corresponding author. Stein Eye Institute, UCLA, 200 Stein Plaza, Los Angeles, CA, 90095, USA.

E-mail address: deng@jsei.ucla.edu (S.X. Deng).

<https://doi.org/10.1016/j.exer.2022.109337>

Received 16 August 2022; Received in revised form 2 November 2022; Accepted 23 November 2022

Available online 23 January 2023

0014-4835/© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

expansion have been developed (Pellegrini, 1997; Rama, 2010). Direct and cultivated LSCs transplantation methods have been reported with high survival rates (Le, 2020), up to 76.6% 10 years after transplantation (Rama, 2010). In the scope of reducing the amount of donor tissue necessary for transplantation, enhancing the outcomes of transplantation from LSCs, and developing targeted therapies, it is important to identify the limbal niche factors involved in LSC regulation. Ultimately, understanding the mechanisms of regulation of the limbal niche will aid in the development of feeder layer-free substrates, mimicking the *in vivo* characteristics of the human limbus, while alleviating any xenobiotic risk (Levis, 2016; Robertson, 2021; Yazdani, 2019; Yazdanpanah, 2019).

Among the multiple and intertwined signaling pathways involved in LSC regulation (Bonnet, 2021b), Wnt signaling pathway activation has been reported in several studies (Han, 2014; Nakatsu, 2013; Vattulainen, 2019). We previously showed that 4 Wnt ligands, WNT2, WNT6, WNT11, and WNT16b, and 4 Wnt inhibitors, dickkopf WNT Signaling Pathway Inhibitor 1 (DKK1), secreted frizzled-related protein 5 (SFRP5), frizzled-related protein (FRZB), and WNT inhibitory factor 1 (WIF1), are preferentially expressed in the limbus compared with the central cornea (Nakatsu, 2011). Activation of the Wnt/ β -catenin pathway by small molecules (Gonzalez, 2019; Nakatsu, 2011; Zhang, 2020) or exogenous Wnt6 (Bonnet, 2021c) can also promote LSC proliferation *in vitro*. Moreover, frizzled7 (Fzd7), the putative stem cell marker and a main Wnt receptor in LSCs, plays a crucial role in the maintenance of the undifferentiated LSC phenotype (Mei, 2014b). This mounting evidence highlights the critical role of Wnt signaling in LSC regulation (Bonnet, 2021a; Bonnet, 2021b).

Wnt proteins are powerful morphogen growth factors with a short range of action (Alexandre, 2014) because of their palmitoylation (Clevers, 2012). This lipid group renders them hydrophobic and thus attracted to neighboring cell membranes or Wnt receptors. Upon secretion of Wnt proteins, the mechanisms of their transport to target cells remain unclear. Different models have been established to support their short range of action (Clevers, 2014). Some Wnt proteins may be incorporated into secretory vesicles that could operate over a short distance, through a paracrine signaling in the stem cell niche. For example in the intestinal niche, Paneth cells secrete Wnt ligands that bind to the intestinal stem cells receptors (Sato, 2011). In addition, interfollicular epidermis stem cells can produce their own Wnt ligands, driving their own self-renewal in an autocrine manner (Lim, 2013). In LSC culture models, the feeder layer, acting as a stem cell niche secreting Wnt ligands (Schrader et al., 2014), is critical for LSC proliferation *in vitro*, suggesting that paracrine signaling occurs in LSC proliferation *in vitro*. Although several transcriptome-profiling studies of the human limbus have been performed, the samples used in the studies contained multiple cell types and did not identify the types of Wnt-producing cells or their precise location (Collin, 2021; Dou, 2021; Kaplan, 2019; Li, 2021; Sun, 2021). Whether LSCs are capable of autocrine Wnt signaling *in vivo* still remains unknown, highlighting the need for more spatio-molecular data, to connect the limbal structure to its function of stem cell renewal.

Traditional histologic methods are limited by the lack of reliable antibodies and the number of markers that are detectable simultaneously; as a result, these methods lack the ability to characterize spatio-molecular details necessary to decipher the complex molecular and cellular structure of the LSC niche. Conversely, the recent development of small-molecule RNA fluorescent *in situ* hybridization (smFISH) methods allows for single-molecule *in situ* RNA localization and quantitation (Femino, 1998; Raj, 2008). Compared with traditional smFISH methods, the RNAscope *in situ* hybridization (ISH) assay could offer higher specificity and sensitivity in different tissue types (Nakajima, 2020; Yu, 2020).

In the present study we evaluated the spatial location of the 4 Wnt ligands and 4 Wnt inhibitors whose expression is upregulated in the limbus (Nakatsu, 2011), and we correlated their location with that of

Fzd7 using the RNAscope ISH technology in the human limbus and central cornea. Wnt signaling pathway molecules were mostly preferentially expressed in the limbal stroma and at the basal epithelial layer of the limbus, and their expression was highly correlated with the localization of Fzd7. A gradient expression pattern was found within limbal epithelial layers. Overall, these results suggest that Wnt signaling in the limbus acts by both paracrine and autocrine signaling, providing new insights into the importance of the LSC niche molecular hierarchical structure to its functions.

2. Material and methods

2.1. Human corneoscleral tissue

Human corneoscleral tissues from 5 healthy donors (median age, 51 years; range, 22–74 years) were obtained with permission for research from Saving Sight (Kansas City, MO, n = 2 donors), EverSight (Clark, NJ, n = 2 donors), and CorneaGen (Orlando, FL, n = 1 donor). Criteria for study eligibility were the following: death-to-preservation time ≤ 8 h (range, 7h10 min to 8h), clear cornea with minimal epithelial exposure and no sloughing, no history of systemic or topical chemotherapy, no history of glaucoma medication use, and no history of ventilation. The use of postmortem human tissues was exempted by the University of California Institutional Review Board (IRB#12–000363). All tissues were preserved at 4 °C in Optisol-GS (Chiron Ophthalmics Inc., Irvine, CA). Under a stereomicroscope on dry ice, tissues were cut into 4 quadrants corresponding to the superior, inferior, nasal, and temporal areas. Each quadrant was embedded in a compound designed for optimal cutting temperature (Tissue-Plus, #4585, Fisher Scientific, Hampton, NH) on dry ice within 30 min of receiving the tissue and preserved at –80 °C. To prevent RNA damage, the tissues from each quadrant were sectioned by a cryostat (ThermoFisher Scientific, Waltham, MA) within a week of cryopreservation, sections were then placed overnight in –80 °C and used the following day. Experiments using human tissues adhered to the tenets of the Declaration of Helsinki.

2.2. *In situ* hybridization

The RNAscope Multiplex Fluorescent V2 Assay (Advanced Cell Diagnostics, Newark, CA) was performed on 12- μ m fresh frozen sections according to the ACD protocol for fresh-frozen tissue without target retrieval (protocol #323100-USM). Each section was contained the limbal region and a section of the peripheral and central cornea. The following modifications to the protocol were made: slides underwent pretreatment for 1 h by baking at 60 °C, permeabilized by using Protease Plus (#322330, ACD, Newark, CA) for 15 min at 40 °C, and fixed in 4% paraformaldehyde (paraformaldehyde 16% aqueous solution, #C004, Electron Microscopy Science, Hatfield, PA, dilution 1:4) for 20 min. Cryosections were then hybridized with 3 mRNA probes per experiment (Supplemental Table 1).

A library of 9 probes, each composed of 20 Z-pairs complementary to sequences of genes involved in Wnt pathway regulation in the limbus was used (Mei, 2014b; Nakatsu, 2011). These probes were WNT2, WNT6, WNT11, WNT16, DKK1, SFRP5, FRZB, WIF1 (Nakatsu, 2011), and a Wnt receptor FZD7 (Mei, 2014b). FZD7 was combined with one Wnt ligand and one Wnt inhibitor, each of different channel, to avoid overlap during signal detection. The probes were amplified according to the manufacturer's instructions and labeled with the fluorophore Opal 520 (Akoya Biosciences, FP1487001KT; dilution ratio, 1:250), Opal 570 (Akoya Biosciences, FP1488001KT; dilution ratio, 1:1500), and Opal 690 (Akoya Biosciences, SKU FP1497001KT; dilution ratio, 1:1500). To confirm the mRNA integrity of tissue blocks, positive-control probes targeted against the human housekeeping genes RNA polymerase II subunit A (POLR2A), peptidylprolyl cis-trans isomerase B (PPIB), and hypoxanthine guanine phosphoribosyltransferase (HPRT), were used on one section in each individual experiment. To confirm signal specificity,

negative-control targeting against bacterial dihydrodipicolinate reductase from *B. subtilis* (*dapB*) gene, scrambled in the 3 channels, was used on one section in each individual experiment. (Supplemental Fig. 1).

2.3. Detection and quantitation of the fluorescence in situ hybridization signals

An upright confocal microscope was used to visualize the fluorescent ISH signals in the limbus and the central cornea at 40X magnification. Images were acquired by using the following parameters: 4 channels (405, 488, 570, 690 nm), sequential mode, 12 bits/pixel, numeric aperture of 0.95, dimension of 1600*640 pixels, and 12 z-stack scans acquired every 1 μm and processed by Maximal Projection on the Olympus FluoView FV1000 v. 4.2.2.9 software.

Quantitation of the spots among the limbal and corneal layers was performed with the surface and spot functions of Imaris software (V. 9.7.0, Imaris, Oxford Instruments, Oxon, UK). Epithelium was defined as the layer of positive nuclei counterstained with DAPI. Segmentation into basal, intermediate, and superficial layers was performed as follow: the basal layer was identified as the layer containing 1 or 2 cells at the junction with the stroma, presenting with very high cell density, having a vertical cell orientation, and possessing a small nucleus when compared with the rest of the epithelium (Romano, 2003). The superficial layer was defined as the most superficial layer of cells, having a larger nucleus and a horizontal cell orientation. The suprabasal layer was defined as the layer between the superficial and basal layers. The corresponding 3 areas were manually drawn and applied onto the 12 z-stacks for volume reconstruction (Supplemental Fig. 2A). Spots were detected by the spot function, which allowed for background subtraction and definition of the spot sizes (i.e., 1.63 μm for Opal 520, 2.01 μm for Opal 570, and 1.61 μm for Opal 647). Spot quality thresholds were set individually for each channel, and the same detection thresholds were used for quantitation consistency among the different experiments. The spots were automatically counted by Imaris processing software (Supplemental Figs. 2B–C).

2.4. Statistical analysis

To control for the thickness variations between limbal and corneal

epithelia, the number of spots counted in each image was normalized to the epithelial volume. Analysis was performed by a biostatistician (C.H. T). Student's *t*-test and analysis of variance were performed using R software (www.r-project.org). A *P* value < 0.05 was considered to indicate statistically significant.

3. Results

3.1. Wnt signaling pathway molecules and LSC marker are differentially expressed in the limbus

Single-molecule RNAscope ISH enabled sensitive and specific in situ transcript detection in the human limbus (Fig. 1A–R) and central cornea (Fig. 1S–T). All 4 Wnt ligands, DKK1, and FZD7 appeared as single spots scattered predominantly in the basal limbal epithelium at a variable density (Fig. 1A–P), whereas SFRP5 was observed among all the limbal epithelial layers (Fig. 1I, M). FRZB and WIF1 were often present as very distinct clusters indicated by high-density signals at the basal layer of the limbus (Fig. 1H,K,L,O,P). These clusters colocalized with areas of FZD7 signals of high density in the basal limbal epithelium (Fig. 2).

Quantitation of these expression profiles retrieved high expression of all 4 Wnt ligands, 4 Wnt inhibitors, and FZD7 in the limbal stroma but not in the central cornea stroma (Supplemental Fig. 3), confirming the role of Wnt signaling in LSC regulation in the limbal niche. All 4 Wnt ligands, 3 inhibitors (WIF1, DKK1, FRZB), and FZD7 were preferentially expressed in the basal layer of the limbus (Fig. 3A–E, K), and the expression decreased in a gradient manner toward the superficial limbal epithelial layer. In the central cornea, expression of 1 Wnt ligand (WNT6), 3 Wnt inhibitors (WIF1, DKK1, SFRP5), and FZD7 was high in the basal layer. The expression level decreased toward the superficial epithelial layer in a gradient manner (Fig. 3F–K).

When compared to the central corneal basal layer, the basal layer of the limbus showed greater spatial expression of 4 Wnt ligands, FRZB, and FZD7 (Fig. 4). No expression in a cluster pattern was detected by any of the probes in the central cornea.




Fig. 1. Expression of Wnt signaling pathways ligands and inhibitors in the human limbus and of Frizzled7 in the human limbus and central cornea. The assay allows specific and sensitive detection of the mRNA expression of Wnt signaling molecules. The mRNA expression of Wnt ligands (A–H) and some Wnt inhibitors (I–P) were preferentially expressed in the basal layer of the limbus. The expression of Fzd7, a putative LSC marker was preferential in the basal layer of the limbus (Q–R). The level of expression was low in the central cornea (S–T). Scale bar: 20 μm . High-magnification pictures represent the white box area in each picture. Scale bar: 5 μm .




Fig. 2. High mRNA level of FRZB and WIF1 co-localized with high level of Fzd7 in the human limbus. FRZB and WIF1 are expressed as clusters colocalizing with high-density Fzd7 signals (white circles). A–C: Fzd7 and FRZB, E–F: Fzd7 and WIF1. Scale bar: 20 μ m.

3.2. The LSC marker Fzd7 is preferentially expressed in the basal layer of the superior quadrant

Spatial expression profiles retrieved preferential expression of FZD7 in the basal layer of the superior quadrant compared to the other limbal quadrants (Fig. 5).

4. Discussion

In this study, we were able to provide a map of the expression patterns of Wnt signaling components previously shown to be preferentially expressed in the limbus by gene profiling. As summarized in Fig. 6, most of the Wnt ligands evaluated are expressed in a gradient manner in the basal limbal epithelial layer decreasing towards the superficial layers, whereas in the central cornea, only Wnt6 presents a decreasing gradient of expression. Conversely, most of the Wnt inhibitors present a decreasing gradient of expression towards the superficial layers in the central cornea, but not in the limbus.

4.1. Expression gradients

Wnt gradients have been reported in many epithelial tissue models such as the hair follicle (Sick, 2006), the intestine crypts (Clevers, 2013), the colon (Gregorieff, 2005), and the adrenal glands (Basham, 2019). Such models provide critical information on the function of some of these factors, where varying levels of Wnt/ β -catenin signaling have different functional consequences for stem cell proliferation and maintenance. In the limbus, the Fzd7, a canonical and non-canonical Wnt receptor (Bonnet, 2021a) was found to co-localize with both Wnt signaling molecules involved in both canonical and non-canonical pathways: Wnt6 is a canonical and non-canonical activator in LSCs (Bonnet, 2021c); Wnt16 is a Wnt/ β -catenin independent activator in LSCs (Zhao, 2022); Wnt2 and Wnt11, act as non-canonical ligands in many cell types (Fan, 2017); and DKK1 is a Wnt/ β -catenin inhibitor (Bourhis, 2011). This observation suggests that both canonical and non-canonical Wnt pathways are tightly regulated by specific stromal and epithelial cells to maintain LSCs homeostasis. Varying temporal and spatial levels of canonical and non-canonical Wnt activation (Kestler, 2011) are likely intertwined and do not function in isolation or

independently. Differential expression pattern between Wnt ligands, which are mostly expressed in the basal layer (Nakatsu, 2011), and Wnt inhibitors, which are more diffusely expressed within the epithelium, may create a highly regulated activation gradient from the basal towards the superficial limbal epithelial layer, similar to that described in the interfollicular epidermidis (Lim, 2013). In this scenario limbal epithelial cell would encounter an increased concentration of Wnt inhibitors as they differentiate and migrate out of the basal layer.

How gradients of secreted Wnt molecules regulate stem cell proliferation and how Wnt molecules travel to interact with their target cells is still unclear. Various modes of Wnt molecules secretion and signaling have been reported, such as extra-cellular vesicles and exosomes (Gross, 2012). For example, miR-6723-5p is upregulated in the limbal basal layers (Ruiz, 2022). miRNAs have also been reported in Wnt signaling regulation during embryogenesis (Nagai, 2019). They are promising therapeutic tools to treat corneal injuries and scars (Deng, 2020a).

Wnt signaling has also been shown to promote corneal wound healing in several animal models by activating metalloproteinase 12 (Lyu, 2005; 2006) and promoting proliferation and migration of corneal epithelial cells (Liang, 2022; Yang, 2020; Zhong, 2021). Understanding the individual molecular mechanisms involved in the limbus and the central cornea could lead to the development of targeted Wnt therapies, promoting both proliferation of LSCs and corneal wound healing.

4.2. Paracrine signaling

The importance of the limbal niche for LSC regulation has been extensively demonstrated over the past 30 years (Dziasko, 2016b; Dziasko, 2014; Levis, 2016; Seyed-Safi, 2020; Tseng, 2016). In ex-vivo LSCs expansion models, the presence of a feeder layer, either in direct contact or using a three-dimensional sandwich model, mimics the niche and supports the cell expansion by secreting soluble niche growth factors, cytokines, and through cell-cell interactions (Mei, 2014a). Moreover, the presence of native limbal stromal cells (Gonzalez, 2013) or of exogenous Wnt activator (Gonzalez, 2019) can increase the expansion efficiency of LSCs in culture. The current manuscript correlates the functions of the limbal stroma and Wnt signaling reported in ex-vivo LSCs proliferation with the preferential expression of Wnt molecules in limbal stromal cells.




Fig. 3. Quantification of Wnt signaling molecules in the limbus and cornea.

A–E: Limbus, F–J: Central cornea Scale bar: 20 μm.

K: Box and whisker plots of the relative level of expression Wnt signaling pathway molecules in the limbus and the central cornea. All four Wnt ligands, WIF1, DKK1, FRZB, and FZD7 are expressed preferentially in the basal limbal epithelial layer, with a significant decrease of expression towards the superficial layer in a gradient manner. In the central cornea, the same gradient of expression is observed for WNT6, WIF1, DKK1, SFRP5, and FZD7. B = basal layer; Sp = suprabasal layer; Su = superficial layer; St = Stroma

Y-axis represents the number of dots detected normalized to the epithelial volume. N = 4 individual experiments performed in duplicates. Supplemental Table 2 details the quantification data.

4.3. Autocrine signaling

The high level of Wnt signaling molecules at the basal layer of the limbal epithelium suggest that LSCs can secrete their own mRNA encoding Wnt ligands and Wnt inhibitors, supporting an autocrine signaling in LSC homeostasis. Autocrine signaling has also been described in interfollicular epidermal stem cells (Lim, 2013), where interfollicular epidermal stem cells expressing *Axin2* can secrete their own Wnt4 and Wnt10a ligands to promote stem cell self-renewal and proliferation. The presence of both autocrine and paracrine signaling highlights the importance of fine tuning Wnt signaling pathway for LSC regulation. These redundant mechanisms (Fig. 6) are critical to avoid aberrant activation of Wnt signaling pathway, preventing ocular surface

neoplasia development.

4.4. *Fzd7* preferential expression in the superior limbus

Fzd7 mRNA expression is higher in the basal epithelial layer of the superior limbus, supporting the role of palisades-rich areas in LSC proliferation. Cultures of LSCs isolated from the superior and inferior limbus have better proliferation capacity than those isolated from the temporal and nasal quadrants (Dziasko, 2014; Meyer-Blazejewska et al., 2010).

The current study has several limitations. First, co-localization is limited to 3 markers due to the number of non-overlapping fluorescent molecules available. In the future, high-definition spatial




Fig. 4. Quantification of mRNA expression profiles of Wnt signaling molecules in the basal epithelial layer of the limbus and central cornea. All 4 Wnt ligands, FRZB, and FZD7 present a preferential expression at the basal epithelial layer of the limbus compared with the basal epithelial layer of the central cornea. Y-axis represents the number of dots detected normalized to the epithelial volume. N = 4 individual experiments performed in duplicates. [Supplemental Table 3](#) details the quantification data.




Fig. 5. The mRNA expression of Fzd7 in the basal limbal epithelial layer is highest in the superior region in the limbus. A- Preferential mRNA expression of FZD7 in the superior quadrant. Scale bar: 20 μ m. B- Box and whisker plots of the relative level of FZD7 single transcripts detected in the basal layer of each quadrant. FZD7 was significantly more expressed in the basal layer of the superior quadrant compared with the three other quadrants ($P < 0.05$). Y-axis represents the number of dots detected normalized to the epithelial volume. N = 4 individual experiments performed in duplicates. [Supplemental Tables 2 and 3](#) detail the quantification data.

transcriptomics (Vickovic, 2019) and slide-seq technologies (Rodrigues, 2019) are expected to improve our spatial understanding of limbal tissue organization. Second, because of the lack of specific antibodies available for the study of the Wnt signaling pathway in the limbus, the findings at the mRNA level could not be confirmed at the protein level. Finally, less than 5% of the limbus was explored, and it would be necessary to perform whole-mounted experiments to obtain a more complete overview when this method becomes feasible.

4.5. Conclusions

In conclusion, several Wnt ligands and inhibitors that are preferentially detected in the limbus compared to the cornea are expressed in a decreasing gradient fashion from the basal limbal epithelial layer and the immediate underlying stroma toward the superficial limbal epithelial layer. As many of these Wnt signaling molecules mediate canonical and/or noncanonical pathways, colocalization of these molecules suggests an interwind and co-dependent relationship of these 2 pathways.




Fig. 6. Model of Wnt signaling expression in the human limbus.

Paracrine and autocrine signaling exist in the human limbus. These redundant mechanisms finely regulate Wnt canonical and non-canonical pathway.

In addition, both autocrine and paracrine effect likely exist in the LSC niche. Taken together, Wnt signaling is tightly controlled in the limbal stem cell niche involving both canonical and noncanonical pathways by a hierarchical expression gradient of Wnt signaling molecules and receptor.

Financial support

This work is supported in part by an unrestricted grant from Research to Prevent Blindness to the Department of Ophthalmology at the University of California, Los Angeles. SXD received grant support from the National Eye Institute (R01 EY021797 and R01 EY028557) and from the California Institute for Regenerative Medicine (TR2-01768, CLIN1-08686, CLIN2-11650) to study limbal stem cells.

Declaration of competing interest

SXD is a consultant for Dompe US, Novartis US, and Claris Biotherapeutics, Inc.; other authors: none.

Data availability

Data will be made available on request.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.exer.2022.109337>.

References

- Alexandre, et al., 2014. Patterning and growth control by membrane-tethered Wingless. *Nature* 505, 180–185. <https://doi.org/10.1038/nature12879>.
- Basham, et al., 2019. A ZNRF3-dependent Wnt/ β -catenin signaling gradient is required for adrenal homeostasis. *Genes Dev.* 33, 209–220. <https://doi.org/10.1101/gad.317412.118>.
- Basu, et al., 2016. Simple limbal epithelial transplantation: long-term clinical outcomes in 125 cases of unilateral chronic ocular surface burns. *Ophthalmology* 123, 1000–1010. <https://doi.org/10.1016/j.ophtha.2015.12.042>.
- Bonnet, et al., 2021a. Wnt signaling activation: targets and therapeutic opportunities for stem cell therapy and regenerative medicine. *RSC Chemical Biology*. <https://doi.org/10.1039/D1CB00063B>.
- Bonnet, et al., 2021b. Human limbal epithelial stem cell regulation, bioengineering and function. *Prog. Retin. Eye Res.*, 100956 <https://doi.org/10.1016/j.preteyeres.2021.100956>.
- Bonnet, et al., 2021c. Wnt6 plays a complex role in maintaining human limbal stem/progenitor cells. *Sci. Rep.* 11, 20948 <https://doi.org/10.1038/s41598-021-00273-y>.
- Bonnet, et al., 2021d. Limbal stem cell diseases. *Exp. Eye Res.*, 108437 <https://doi.org/10.1016/j.exer.2021.108437>.
- Bourhis, et al., 2011. Wnt antagonists bind through a short peptide to the first β -propeller domain of LRP5/6. *Structure* 19, 1433–1442. <https://doi.org/10.1016/j.str.2011.07.005>.
- Clevers, 2013. The intestinal crypt, a prototype stem cell compartment. *Cell* 154, 274–284. <https://doi.org/10.1016/j.cell.2013.07.004>.
- Clevers, et al., 2014. Stem cell signaling. An integral program for tissue renewal and regeneration: Wnt signaling and stem cell control. *Science (New York, N.Y.)* 346. <https://doi.org/10.1126/science.1248012>.
- Clevers, Nusse, 2012. Wnt/ β -catenin signaling and disease. *Cell* 149, 1192–1205. <https://doi.org/10.1016/j.cell.2012.05.012>.
- Collin, et al., 2021. A single cell atlas of human cornea that defines its development, limbal progenitor cells and their interactions with the immune cells. *Ocul. Surf.* <https://doi.org/10.1016/j.jtos.2021.03.010>.
- Cotsarelis, et al., 1989. Existence of slow-cycling limbal epithelial basal cells that can be preferentially stimulated to proliferate: implications on epithelial stem cells. *Cell* 57, 201–209. [https://doi.org/10.1016/0092-8674\(89\)90958-6](https://doi.org/10.1016/0092-8674(89)90958-6).
- Davanger, Evensen, 1971. Role of the pericorneal papillary structure in renewal of corneal epithelium. *Nature* 229, 560–561. <https://doi.org/10.1038/229560a0>.
- Deng, et al., 2019. Global consensus on definition, classification, diagnosis, and staging of limbal stem cell deficiency. *Cornea* 38, 364–375. <https://doi.org/10.1097/ico.0000000000001820>.
- Deng, et al., 2020a. Therapeutic potential of extracellular vesicles for the treatment of corneal injuries and scars. *Transl Vis Sci Technol* 9, 1. <https://doi.org/10.1167/tvst.9.12.1>.
- Deng, et al., 2020b. Global consensus on the management of limbal stem cell deficiency. *Cornea* 39, 1291–1302. <https://doi.org/10.1097/ico.0000000000002358>.
- Dou, et al., 2021. Molecular identity of human limbal heterogeneity involved in corneal homeostasis and privilege. *Ocul. Surf.* <https://doi.org/10.1016/j.jtos.2021.04.010>.
- Dziasko, et al., 2014. Localisation of epithelial cells capable of holoclone formation in vitro and direct interaction with stromal cells in the native human limbal crypt. *PLoS One* 9, e94283. <https://doi.org/10.1371/journal.pone.0094283>.
- Dziasko, Daniels, 2016a. Anatomical features and cell-cell interactions in the human limbal epithelial stem cell niche. *Ocul. Surf.* 14, 322–330. <https://doi.org/10.1016/j.jtos.2016.04.002>.
- Dziasko, Daniels, 2016b. Anatomical features and cell-cell interactions in the human limbal epithelial stem cell niche. *Ocul. Surf.* 14 <https://doi.org/10.1016/j.jtos.2016.04.002>.
- Fan, et al., 2017. Noncanonical Wnt signaling plays an important role in modulating canonical Wnt-regulated stemness, proliferation and terminal differentiation of hepatic progenitors. *Oncotarget* 8, 27105–27119. <https://doi.org/10.18632/oncotarget.15637>.
- Femino, et al., 1998. Visualization of single RNA transcripts in situ. *Science* 280, 585–590. <https://doi.org/10.1126/science.280.5363.585>.
- Gonzalez, Deng, 2013. Presence of native limbal stromal cells increases the expansion efficiency of limbal stem/progenitor cells in culture. *Exp. Eye Res.* 116, 169–176. <https://doi.org/10.1016/j.exer.2013.08.020>.
- Gonzalez, et al., 2019. Wnt signaling is required for the maintenance of human limbal stem/progenitor cells in vitro. *Invest. Ophthalmol. Vis. Sci.* 60, 107–112. <https://doi.org/10.1167/iov.18-25740>.
- Gregorieff, et al., 2005. Expression pattern of Wnt signaling components in the adult intestine. *Gastroenterology* 129, 626–638. <https://doi.org/10.1016/j.gastro.2005.06.007>.
- Gross, et al., 2012. Active Wnt proteins are secreted on exosomes. *Nat. Cell Biol.* 14, 1036–1045. <https://doi.org/10.1038/ncb2574>.
- Han, et al., 2014. Integration of BMP/Wnt signaling to control clonal growth of limbal epithelial progenitor cells by niche cells. *Stem Cell Res.* 12, 562–573. <https://doi.org/10.1016/j.scr.2014.01.003>.
- Kaplan, et al., 2019. Single-cell RNA transcriptome helps define the limbal/corneal epithelial stem/early transit amplifying cells and how autophagy affects this population. *Invest. Ophthalmol. Vis. Sci.* 60, 3570–3583. <https://doi.org/10.1167/iov.19-27656>.
- Kestler, Kühl, 2011. Generating a Wnt switch: it's all about the right dosage. *J. Cell Biol.* 193, 431–433. <https://doi.org/10.1083/jcb.201103167>.
- Le, et al., 2020. Outcomes of limbal stem cell transplant: a meta-analysis. *JAMA ophthalmology* 138, 660–670. <https://doi.org/10.1001/jamaophthalmol.2020.1120>.
- Levis, Daniels, 2016. Recreating the human limbal epithelial stem cell niche with bioengineered limbal crypts. *Curr. Eye Res.* 41, 1153–1160. <https://doi.org/10.3109/02713683.2015.1095932>.
- Li, et al., 2021. Single-cell transcriptomics identifies limbal stem cell population and cell types mapping its differentiation trajectory in limbal basal epithelium of human cornea. *Ocul. Surf.* 20, 20–32. <https://doi.org/10.1016/j.jtos.2020.12.004>.
- Liang, et al., 2022. Pathogenic role of diabetes-induced overexpression of kallistatin in corneal wound healing deficiency through inhibition of canonical Wnt signaling. *Diabetes* 71, 747–761. <https://doi.org/10.2337/db21-0740>.

- Lim, et al., 2013. Interfollicular epidermal stem cells self-renew via autocrine Wnt signaling. *Science* 342, 1226–1230. <https://doi.org/10.1126/science.1239730>.
- Lyu, Joo, 2005. Wnt-7a up-regulates matrix metalloproteinase-12 expression and promotes cell proliferation in corneal epithelial cells during wound healing. *J. Biol. Chem.* 280, 21653–21660. <https://doi.org/10.1074/jbc.M500374200>.
- Lyu, Joo, 2006. Expression of Wnt and MMP in epithelial cells during corneal wound healing. *Cornea* 25, S24–S28. <https://doi.org/10.1097/01.ico.0000247209.01262.4e>.
- Mei, et al., 2014a. A three-dimensional culture method to expand limbal stem/progenitor cells. *Tissue Eng.* 20, 393–400. <https://doi.org/10.1089/ten.TEC.2013.0246>.
- Mei, et al., 2014b. Frizzled 7 maintains the undifferentiated state of human limbal stem/progenitor cells. *Stem Cell.* 32, 938–945. <https://doi.org/10.1002/stem.1582>.
- Meyer-Blazejewska, et al., 2010. Preservation of the limbal stem cell phenotype by appropriate culture techniques. *Invest. Ophthalmol. Vis. Sci.* 51, 765–774. <https://doi.org/10.1167/iovs.09-4109>.
- Nagai, et al., 2019. MicroRNAs control eyelid development through regulating Wnt signaling. *Dev. Dynam.* 248, 201–210. <https://doi.org/10.1002/dvdy.10>.
- Nakajima, et al., 2020. Characterization of LGR5 expression in poorly differentiated colorectal carcinoma with mismatch repair protein deficiency. *BMC Cancer* 20, 319. <https://doi.org/10.1186/s12885-020-06791-8>.
- Nakatsu, et al., 2011. Wnt/beta-catenin signaling regulates proliferation of human cornea epithelial stem/progenitor cells. *Invest. Ophthalmol. Vis. Sci.* 52, 4734–4741. <https://doi.org/10.1167/iovs.10-6486>.
- Nakatsu, et al., 2013. Preferential biological processes in the human limbus by differential gene profiling. *PLoS One* 8, e61833. <https://doi.org/10.1371/journal.pone.0061833>.
- Pellegrini, et al., 1997. Long-term restoration of damaged corneal surfaces with autologous cultivated corneal epithelium. *Lancet* 349, 990–993. [https://doi.org/10.1016/s0140-6736\(96\)11188-0](https://doi.org/10.1016/s0140-6736(96)11188-0).
- Raj, et al., 2008. Imaging individual mRNA molecules using multiple singly labeled probes. *Nat. Methods* 5, 877–879. <https://doi.org/10.1038/nmeth.1253>.
- Rama, et al., 2010. Limbal stem-cell therapy and long-term corneal regeneration. *N. Engl. J. Med.* 363, 147–155. <https://doi.org/10.1056/NEJMoa0905955>.
- Robertson, et al., 2021. Regulation of limbal epithelial stem cells: importance of the niche. *Int. J. Mol. Sci.* 22, 11975.
- Rodrigues, et al., 2019. Slide-seq: a scalable technology for measuring genome-wide expression at high spatial resolution. *Science* 363, 1463–1467. <https://doi.org/10.1126/science.aaw1219>.
- Romano, et al., 2003. Different cell sizes in human limbal and central corneal basal epithelia measured by confocal microscopy and flow cytometry. *Invest. Ophthalmol. Vis. Sci.* 44, 5125–5129. <https://doi.org/10.1167/iovs.03-0628>.
- Ruiz, et al., 2022. Extracellular miR-6723-5p could serve as a biomarker of limbal epithelial stem/progenitor cell population. *Biomark Res* 10, 36. <https://doi.org/10.1186/s40364-022-00384-2>.
- Sato, et al., 2011. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature* 469, 415–418. <https://doi.org/10.1038/nature09637>.
- Scadden, 2006. The stem-cell niche as an entity of action. *Nature* 441, 1075–1079. <https://doi.org/10.1038/nature04957>.
- Schrader, et al., 2014. Wnt signalling in an in vitro niche model for conjunctival progenitor cells. *J Tissue Eng Regen Med* 8, 969–977. <https://doi.org/10.1002/term.1599>.
- Seyed-Safi, Daniels, 2020. The limbus: structure and function. *Exp. Eye Res.* 197, 108074. <https://doi.org/10.1016/j.exer.2020.108074>.
- Sick, et al., 2006. WNT and DKK determine hair follicle spacing through a reaction-diffusion mechanism. *Science* 314, 1447–1450. <https://doi.org/10.1126/science.1130088>.
- Sun, et al., 2021. Time-course single-cell RNA sequencing reveals transcriptional dynamics and heterogeneity of limbal stem cells derived from human pluripotent stem cells. *Cell Biosci.* 11, 24. <https://doi.org/10.1186/s13578-021-00541-4>.
- Tseng, 1989. Concept and application of limbal stem cells. *Eye* 3 (Pt 2), 141–157. <https://doi.org/10.1038/eye.1989.22>.
- Tseng, et al., 2016. Niche regulation of limbal epithelial stem cells: relationship between inflammation and regeneration. *Ocul. Surf.* 14, 100–112. <https://doi.org/10.1016/j.jtos.2015.12.002>.
- Vattulainen, et al., 2019. Modulation of Wnt/BMP pathways during corneal differentiation of hPSC maintains ABCG2-positive LSC population that demonstrates increased regenerative potential. *Stem Cell Res. Ther.* 10, 236. <https://doi.org/10.1186/s13287-019-1354-2>.
- Vickovic, et al., 2019. High-definition spatial transcriptomics for in situ tissue profiling. *Nat. Methods* 16, 987–990. <https://doi.org/10.1038/s41592-019-0548-y>.
- Yang, et al., 2020. Insulin promotes corneal nerve repair and wound healing in type 1 diabetic mice by enhancing wnt/ β -catenin signaling. *Am. J. Pathol.* 190, 2237–2250. <https://doi.org/10.1016/j.ajpath.2020.08.006>.
- Yazdani, et al., 2019. A hyaluronan hydrogel scaffold for culture of human oral mucosal epithelial cells in limbal stem-cell therapy. *Bioengineering (Basel)* 6. <https://doi.org/10.3390/bioengineering6040097>.
- Yazdanpanah, et al., 2019. Strategies for reconstructing the limbal stem cell niche. *Ocul. Surf.* 17, 230–240. <https://doi.org/10.1016/j.jtos.2019.01.002>.
- Yu, et al., 2020. Epithelial Wnt10a is essential for tooth root furcation morphogenesis. *J. Dent. Res.* 99, 311–319. <https://doi.org/10.1177/0022034519897607>.
- Zhang, et al., 2020. A small-molecule Wnt mimic improves human limbal stem cell ex vivo expansion. *iScience* 23, 101075. <https://doi.org/10.1016/j.isci.2020.101075>.
- Zhao, et al., 2022. WNT16B enhances the proliferation and self-renewal of limbal epithelial cells via CXCR4/MEK/ERK signaling. *Stem Cell Rep.* <https://doi.org/10.1016/j.stemcr.2022.03.001>.
- Zhong, et al., 2021. Ocular Wnt/ β -catenin pathway inhibitor XAV939-loaded liposomes for treating alkali-burned corneal wound and neovascularization. *Front. Bioeng. Biotechnol.* 9, 753879. <https://doi.org/10.3389/fbioe.2021.753879>.