



HAL
open science

Should initial ICU admission become a standard of care for acute bacterial meningitis ?

Michael Thy, Claire Dupuis, Jean-François Timsit, Romain Sonnevile

► To cite this version:

Michael Thy, Claire Dupuis, Jean-François Timsit, Romain Sonnevile. Should initial ICU admission become a standard of care for acute bacterial meningitis?. *Annals of Intensive Care*, 2024, 14 (1), pp.48. 10.1186/s13613-024-01277-3 . hal-04591976

HAL Id: hal-04591976

<https://u-paris.hal.science/hal-04591976>

Submitted on 29 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EDITORIAL

Open Access



Should initial ICU admission become a standard of care for acute bacterial meningitis ?

Michael Thy^{1,2*} , Claire Dupuis³, Jean-François Timsit^{1,4} and Romain Sonnevill^{1,4}

The decision to admit adult patients with bacterial meningitis to the intensive care unit (ICU) is usually based on severe illness at hospital presentation, including altered mental status, seizures, respiratory and/or cardiovascular complications. However, as some patients may deteriorate rapidly within the first hours after hospital presentation, prophylactic ICU admission could be discussed in stable patients at high risk of secondary deterioration. However, the profile of these patients is not known. Two recent studies published in “Annals of Intensive Care” provide important information on this topic [1, 2]. A summary and comparison of the patients directly admitted to the ICU from the two studies are shown in Table 1.

The first study is the MeninGene nationwide study, conducted from 2006 to 2022 in the Netherlands, which presents a comprehensive examination of adults with community-acquired bacterial meningitis who eventually required ICU admission, either directly or secondarily [1]. This study provides precise estimates of intensive care needs in this population, with 51% of patients requiring initial ICU admission. This study confirms that the prognosis of such patients remains poor

with high unfavorable outcome (47%) and hospital mortality (22%) rates. Factors independently associated with ICU admission were age, male gender, immunocompromised state, and factors associated with pneumococcal etiology. These factors may help to identify high risk patients and assist clinicians with complex decisions of ICU admission. Of particular concern, were the subgroup of patients initially admitted to non-ICU wards but subsequently transferred to the ICU, constituting 15% of cases. Importantly, this subgroup faced even higher rates of unfavorable outcome (66%) and mortality (30%) than those of directly admitted patients.

The second study based is a retrospective analysis of the French medico-administrative database on 4052 cases of pneumococcal meningitis with sepsis criteria recruited from 2011 to 2020 who eventually received care in the ICU [2]. Among them, 50% were directly admitted to the ICU and the remaining 50% were secondarily admitted. After adjustment for confounders (including gender, cardiovascular comorbidities, chronic obstructive pulmonary disease, diabetes, chronic kidney disease, neurological, respiratory and hematological failures on admission, associated diagnosis of endocarditis), direct ICU admission was associated with lower hospital mortality rates when compared to delayed admission (adjusted odds ratio 0.67, 95%CI 0.56–0.80). Although the reason for secondary ICU admission in both studies remains speculative, it is likely that these patients experienced early deterioration because of complications of neurologic or systemic origin. Adding the recent data from EURECA study and a recent large retrospective study [3, 4], we suggest a management algorithm of

*Correspondence:

Michael Thy
michael245thy@gmail.com

¹Medical and Infectious Diseases ICU, Bichat Claude Bernard University Hospital, Université Paris Cité, AP-HP, Paris, France

²EA7323, Pharmacology and Drug Evaluation in Children and Pregnant Women, Université Paris Cité, Paris, France

³Department of Intensive Care Medicine, Gabriel-Montpied University Hospital, Clermont- Ferrand, France

⁴Decision Sciences in Infectious Diseases control and care, UMR 1137-IAME Team:INSERM Université Paris Cité, Paris 75018, France



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Table 1 Comparison of the two studies for bacterial meningitis initially or directly admitted to the ICU

Direct ICU admission for bacterial meningitis	Chekrouni et al. [1]	Thy et al. [2]
Design	Prospective cohort of community-acquired bacterial meningitis (MeninGene study)	Retrospective analysis of pneumococcal meningitis with sepsis (PMSI database)
Country	Netherlands	France
Date of inclusion	From 2006 to 2022	From 2011 to 2020
Main characteristics	<i>N</i> = 2709	<i>N</i> = 4052
Initial/direct ICU admission rate	1369 (51%)	2006 (50%)
Median age (IQR), years	61 (49–69)	60 (49–70)
Male sex	737/1369 (54%)	1141 (57%)
<i>S. pneumoniae</i>	1071/1369 (78%)	100%
Associated infection		
Endocarditis	26/1312 (2%)	139 (7%)
Pneumonia	141/1313 (11%)	712 (36%)
Complications		
Coma (GCS < 8)	431/1360 (32%)	1117 (56%)
Focal neurological deficits	362 of 1220 (30%)	373 (19%)
Seizures	230/1313 (18%)	231 (12%)
Cranial nerve palsy / Brainstem failure	97/1074 (9%)	356 (18%)
Vascular complications*	244 (18%)	150 (8%)
Hydrocephalus	72/1241 (6%)	27 (1%)
Organ failure on admission		
Cardiovascular failure on admission	220/1272 (17%)	1188 (59%)
Respiratory failure on admission	470/1300 (36%)	1437 (72%)
Outcomes		
In-hospital mortality	304 (22%)	533 (27%)
Unfavorable outcome (GOS 2–4 or discharge to readaptation)	339 (25%)	337 (17%)
Median ICU length of stay (IQR), days	3 (2–8)	7 (3–15)
Median hospital length of stay (IQR), days	15 (12–24)	21 (13–37)

ICU: intensive or intermediate care unit, PMSI: Programme de Médicalisation des Systèmes d'Information, GCS: Glasgow coma scale, GOS: Glasgow outcome scale

*Vascular complications included ischemic stroke, transient ischemic stroke and cerebral venous thrombosis

intensive care unit (ICU) admission for suspected or confirmed acute bacterial meningitis on Fig. 1.

The implications of these two multicenter studies are profound as they help to identify patients at risk of poor outcome and to highlight the potential benefit of a more aggressive approach toward direct ICU admission, particularly for pneumococcal cases with sepsis. Indeed, such findings underscore the dynamic nature of patient trajectories and the urgency of refining admission protocols to ensure timely and appropriate care.




Fig. 1 Management of intensive care unit (ICU) admission for suspected or confirmed acute bacterial meningitis. Legends: *Altered mental status, brain-stem involvement/cranial nerve dysfunction and/or seizures. **With early discharge to medical wards after improvement

Acknowledgements

The authors wish to thank all the patients and all the healthcare workers of French hospitals.

Author contributions

MT, CD, RS and JFT drafted the manuscript and critically revised the manuscript.

Funding

The study was not externally funded.

Data availability

Data available on request.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

All authors read, revised and approved the final manuscript for publication.

Competing interests

Not applicable.

References

1. Chekrouni N, Kroon M, Drost EHGM, van Soest TM, Bijlsma MW, Brouwer MC, et al. Characteristics and prognostic factors of bacterial meningitis in the intensive care unit: a prospective nationwide cohort study. *Ann Intensive Care*. 2023;13:124.
2. Thy M, Dupuis C, Mageau A, Mourvillier B, Bouadma L, Ruckly S, et al. Impact of direct ICU admission of pneumococcal meningitis in France: a retrospective analysis of a French medico-administrative (PMSI) database. *Ann Intensive Care*. 2024;14:15.
3. Sonnevile R, de Montmollin E, Contou D, Ferrer R, Gurjar M, Klouche K et al. Clinical features, etiologies, and outcomes in adult patients with meningococcal meningitis requiring intensive care (EURECA): an international prospective multicenter cohort study. *Intensive Care Med* [Internet]. 2023 [cited 2023 Apr 7]; <https://doi.org/10.1007/s00134-023-07032-9>.
4. Donovan J, Glover A, Gregson J, Hitchings AW, Wall EC, Heyderman RS. A retrospective analysis of 20,178 adult neurological infection admissions to United Kingdom critical care units from 2001 to 2020. *BMC Infect Dis*. 2024;24:132.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 3 February 2024 / Accepted: 11 March 2024

Published online: 01 April 2024